

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 471/04, 498/04, A61K 31/55	A1	(11) International Publication Number: WO 99/06403 (43) International Publication Date: 11 February 1999 (11.02.99)
(21) International Application Number: PCT/US98/15487 (22) International Filing Date: 24 July 1998 (24.07.98) (30) Priority Data: 08/903,370 30 July 1997 (30.07.97) US 08/941,707 30 July 1997 (30.07.97) US (71) Applicant: AMERICAN HOME PRODUCTS CORPORATION [US/US]; Five Giralda Farms, Madison, NJ 07940-0874 (US). (72) Inventors: FAILLI, Amedeo, Arturo; 14 Landing Lane, Princeton Junction, NJ 08550 (US). SHUMSKY, Jay, Scott; Princeton Arms Apartments, #81 South, Cranbury, NJ 08512 (US). STEFFAN, Robert, John; 263 Wheatsheaf Lane, Langhorne, PA 19047 (US). (74) Agents: ALICE, Ronald, W.; American Home Products Corporation, Patent Law Dept. -2B, One Campus Drive, Parsippany, NJ 07054 (US) et al.		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: TRICYCLIC VASOPRESSIN AGONISTS <div data-bbox="467 1167 959 1499"><p style="text-align: right;">(I)</p></div> (57) Abstract <p>The present invention provides compounds of general formula (I): wherein W is O or NH, optionally substituted, as well as methods and pharmaceutical compositions utilising these compounds for the treatment of disorder which may be remedied or alleviated by vasopressin agonist activity, including diabetes insipidus, nocturnal enuresis, nocturia, urinary incontinence, bleeding and coagulation disorders, or temporary delay of urination.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TRICYCLIC VASOPRESSIN AGONISTS

This invention concerns tricyclic pyrido compounds, or pharmaceutical salts thereof, which act as vasopressin agonists, as well as methods of treatment and pharmaceutical compositions utilizing the compounds.

Background of the Invention

Vasopressin (antidiuretic hormone, ADH) a non-peptide hormone and neurotransmitter, is synthesized in the supraoptic nuclei of the hypothalamus of the brain and is transported through the supraoptico-hypophyseal tract to the posterior pituitary where it is stored. Upon sensing an increase in plasma osmolality by brain osmoreceptors or a decrease in blood volume or blood pressure (detected by the baroreceptors and volume receptors), vasopressin is released into the blood circulation and it activates V_{1a} receptors on blood vessels causing vasoconstriction to raise blood pressure and vasopressin V_2 receptors of the nephron of the kidney causing retention mainly of water, and to a lesser degree electrolytes, to expand the blood volume (Cervoni and Chan, *Diuretic Agents*, in Kirk-Othmer, Encyclopedia of Chemical Technology, 4th ed., Wiley, volume 8, 398-432, (1993)). The existence of vasopressin in the pituitary was known as early as 1895 (Oliver and Schaefer, *J. Physiol. (London)*, **18**, 277-279, (1895)). The determination of the structure and the total synthesis of vasopressin were accomplished by du Vigneaud and coworkers in 1954 (du Vigneaud, Gish and Katsoyannis, *J. Am. Chem. Soc.*, **76**, 4751-4752, (1954)).

The actions of vasopressin V_{1a} receptors are mediated through the phosphatidylinositol pathway. Activation of vasopressin V_{1a} receptors causes contraction of the smooth muscle of the blood vessels to raise blood pressure. The actions of the vasopressin V_2 receptors are mediated through activation of the adenylate cyclase system and elevation of intracellular levels of cAMP. The activation of vasopressin V_2 receptors by vasopressin or vasopressin-like (peptidic or non-peptidic) compounds increases water permeability of the collecting ducts of the nephron and permits the reabsorption of a large quantity of free water. The end result is the formation and excretion of a concentrated urine, with a decrease in urine volume and an increase in urinary osmolality.

Vasopressin plays a vital role in the conservation of water by concentrating the urine at the site of the collecting ducts of the kidney. The collecting ducts of the kidney are relatively impermeable to water without the presence of vasopressin at the receptors and therefore, the hypotonic fluid formed after filtering through the glomeruli, passing the proximal convoluted tubule, the loops of Henle, and the distal convoluted tubules, will be excreted as dilute urine. However, during dehydration, volume depletion or blood loss, vasopressin is released from the brain and activates the vasopressin V_2 receptors in the collecting ducts of the kidney rendering the ducts very permeable to water; hence water is reabsorbed and a concentrated urine is excreted. In patients and animals with central or neurogenic diabetes insipidus, the synthesis of vasopressin in the brain is defective and therefore, they produce no or very little vasopressin, but their vasopressin receptors in the kidneys are normal. Because they cannot concentrate the urine, they may produce as much as 10 times the urine volumes of their healthy counterparts and they are very sensitive to the action of vasopressin and vasopressin V_2 agonists. Vasopressin and desmopressin, which is a peptide analog of the natural vasopressin, are being used in patients with central diabetes insipidus. Vasopressin V_2 agonists are also useful for the treatment of nocturnal enuresis, nocturia, urinary incontinence and temporary delay of urination whenever desirable.

Vasopressin, through activation of its V_{1a} receptors, exerts vasoconstricting effects so as to raise blood pressure. A vasopressin V_{1a} receptor antagonist will counteract this effect. Vasopressin and vasopressin agonists release factor VIII and von Willebrand factor so they are useful for the treatment of bleeding disorders, such as hemophilia. Vasopressin and vasopressin-like agonists also release tissue-type plasminogen activator (t-PA) into the blood circulation so they are useful in dissolving blood clots such as in patients with myocardial infarction and other thromboembolic disorders (Jackson, "Vasopressin and other agents affecting the renal conservation of water", in Goodman and Gilman, *The Pharmacological Basis of Therapeutics*, 9th ed., Hadman, Limbird, Molinoff, Ruddon and Gilman Eds., McGraw-Hill, New York, pp. 715-731 (1996); Lethagen, *Ann. Hematol.* **69**, 173-180 (1994); Cash et al., *Brit. J. Haematol.*, **27**, 363-364 (1974); David, *Regulatory Peptides*, **45**, 311-317 (1993); Burggraaf et al., *Cli. Sci.*, **86**, 497-503 (1994)).

The following prior art references describe peptidic vasopressin antagonists: Manning et al., *J. Med. Chem.*, **35**, 382 (1992); Manning et al., *J. Med. Chem.*, **35**, 3895 (1992); Gavras and Lammek, U.S. Patent 5,070,187 (1991); Manning and Sawyer, U.S. Patent 5,055,448 (1991); Ali, U.S. Patent 4,766,108 (1988); Ruffolo et al., *Drug News and Perspectives* **4**(4), 217 (May 1991). Williams et al., have reported on potent hexapeptide oxytocin antagonists [*J. Med. Chem.*, **35**, 3905 (1992)] which also exhibit weak vasopressin antagonistic activity in binding to V₁ and V₂ receptors. Peptidic vasopressin antagonists suffer from a lack of oral activity and many of these peptides are non-selective antagonists since they also exhibit partial agonist activity.


Non-peptidic vasopressin antagonists have recently been disclosed. Albright et al. describe tricyclic diazepines as vasopressin and oxytocin antagonists in U.S. Patent 5,516,774 (1996); tetrahydrobenzodiazepine derivatives as vasopressin antagonists are disclosed in J.P. 0801460-A (1996); Ogawa et al., disclose benzoheterocyclic derivatives as vasopressin and oxytocin antagonists, and as vasopressin agonists in WO 9534540-A; Albright et al., disclose tricyclic benzazepine derivatives as vasopressin antagonists in U.S. Patent 5,512,563 (1996); and Venkatesan et al., disclose tricyclic benzazepine derivatives as vasopressin and oxytocin antagonists in U.S. Patent 5,521,173 (1996).

As mentioned above, desmopressin (1-desamino-8-D-arginine vasopressin) (Huguenin and Boissonnas, *Helv. Chim. Acta*, **49**, 695 (1966)) is a vasopressin agonist. The compound is a synthetic peptide with variable bioavailability. An intranasal route is poorly tolerated and an oral formulation for nocturnal enuresis requires a 10-20 fold greater dose than the intranasal administration.

The compounds of this invention are non-peptidic and have a good oral bioavailability. They are selective vasopressin V₂ agonists, and have no V_{1a} agonist effects so they do not raise blood pressure. In contrast, the prior art compounds (except some in WO 9534540-A) are vasopressin antagonists.

SUMMARY OF THE INVENTION

This invention relates to novel compounds selected from those of Formula (I):



(I)


wherein:

W is O, or NR⁶

5 A and B are, independently, carbon or nitrogen;

R¹ is $\text{—C}\equiv\text{C—R}^9$, alkanoyl of 2 to 7 carbon atoms or a moiety selected from the group of:

-5-



R^2 , R^3 and R^5 are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or perfluoroalkyl of 1 to 6 carbons;


R^4 is hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, alkoxyalkyl of 2 to 7 carbon atoms, optionally substituted aralkyl of 7 to 12 carbon atoms, or an acyl substituent selected from the group consisting of alkanoyl of 2 to 7 carbon atoms, alkenoyl of 3 to 7 carbon atoms, cycloalkanoyl of 3 to 7 carbon atoms, arylalkanoyl having an alkane chain of from 1 to 6 carbon atoms, aroyl or heteroaroyl of 7 to 13 carbon atoms;

R^6 is hydrogen, acyl of 2 to 6 carbon atoms, straight chain alkyl of 1 to 6 carbon atoms, or branched chain alkyl of 3 to 7 carbon atoms;

X and Y are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, alkoxyalkyl of 2 to 7 carbon atoms, halogen, straight or branched chain alkoxy
 5 of 1 to 6 carbon atoms, hydroxy, CF_3 , or perfluoroalkyl of 2 to 6 carbons;

Z is hydrogen or a straight chain alkyl group of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, halogen,
 alkoxyalkyl of 2 to 7 carbons, or hydroxyalkyl of 1 to 6 carbons, or
 10 $\text{CH}_2\text{NR}^7\text{R}^8$;

R^7 and R^8 are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, aryl, or arylalkyl; or taken together with the nitrogen they form a five or six membered ring optionally
 15 containing one or more additional heteroatoms, preferably selected from N, S or O, such as



20 R^9 is independently, hydrogen, a silyl containing group, or a lower alkyl of 1 to 6 carbons; and

R^{10} is a straight chain alkyl of 1 to 6 carbon atoms;

or a pharmaceutically acceptable salt, ester or prodrug form thereof.

25

R^1 is preferably a moiety selected from:

- group a defined above wherein two of R^2 , R^3 and R^5 are hydrogen and the third is a straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3

to 7 carbon atoms or perfluoroalkyl of 1 to 6 carbons, more preferably where the third is selected from methyl, t-butyl and trifluoromethyl,

5 - group f defined above wherein one of R^2 and R^4 is hydrogen and the other is a straight chain alkyl of 1 to 6 carbon atoms or branched chain alkyl of 3 to 7 carbon atoms, more preferably it is methyl,

10 - group e or i defined above wherein R^4 is hydrogen and R^2 is a straight chain alkyl of 1 to 6 carbon atoms or branched chain alkyl of 3 to 7 carbon atoms, more preferably it is methyl.

15 R^2 , R^3 and R^5 are each preferably hydrogen, a straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms or perfluoroalkyl of 1 to 6 carbons, more preferably hydrogen, methyl, t-butyl or trifluoromethyl. Preferred embodiments are those wherein two of R^2 , R^3 and R^5 are hydrogen and the third is a straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms or perfluoroalkyl of 1 to 6 carbons, more preferably where the third is selected from methyl, t-butyl and trifluoromethyl.

20 R^4 is preferably hydrogen, a straight chain alkyl of 1 to 6 carbon atoms or a branched chain alkyl of 3 to 7 carbon atoms, more preferably hydrogen or methyl.

25 R^6 is preferably hydrogen or a straight chain alkyl of 1 to 6 carbon atoms, more preferably hydrogen or methyl.



30 X and Y are each preferably hydrogen, halogen, perfluoroalkyl of 2 to 6 carbons, straight chain alkyl of 1 to 6 carbon atoms or a branched chain alkyl of 3 to 7 carbon atoms, more preferably hydrogen, chloro, bromo, fluoro, trifluoromethyl, or methyl. Preferably at least one of X and Y is hydrogen, more preferably X is hydrogen.

35 Z is preferably hydrogen.

R^7 and R^8 are each preferably hydrogen or a straight chain alkyl of 1 to 6 carbon atoms, more preferably hydrogen or methyl.


Within the scope of this invention are subgeneric groups of the abovementioned compounds wherein W is O and A, B, X, Y, Z and R^1 - R^{10} are as defined above; and W is NR^6 and A, B, X, Y, Z and R^1 - R^{10} are as defined above.

- 5 Among the more preferred compounds of this invention are those selected from Formula (Ia) or (Ib):



- 10 wherein:

R^1 is alkanoyl of 2 to 7 carbon atoms or a moiety selected from the group of:



R^2 , R^3 and R^5 are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or perfluoroalkyl of 1 to 6 carbons;

R^4 is hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, alkoxyalkyl of 2 to 7 carbon atoms, optionally substituted aralkyl of 7 to 12 carbon atoms, or an acyl substituent selected from the group consisting of alkanoyl of 2 to 7 carbon atoms, alkenoyl of 3 to 7 carbon atoms, cycloalkanoyl of 3 to 7 carbon atoms, aroyl or heteroaroyl of 7 to 12 carbon atoms, optionally substituted with one or two alkyl groups of 1 to 6 carbon atoms, or arylalkanoyl of 1 to 6 carbon atoms;


R^6 is hydrogen, acyl of 2 to 6 carbon atoms, straight chain alkyl of 1 to 6 carbon atoms, or branched chain alkyl of 3 to 7 carbon atoms;

X and Y are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, alkoxyalkyl of 2 to 7 carbon atoms, halogen (including chlorine, bromine, fluorine, and iodine), straight chain or branched chain alkoxy of 1 to 6 carbons, hydroxy, CF_3 , or perfluoroalkyl of 2 to 6 carbons;

Z is hydrogen or a straight chain alkyl group of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, halogen, alkoxyalkyl of 2 to 7 carbons, hydroxyalkyl of 1 to 6 carbons, or $\text{CH}_2\text{NR}^7\text{R}^8$;

- 5 R^7 and R^8 are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, aryl or arylalkyl; or taken together with the nitrogen they form a five or six membered ring optionally containing one or more additional heteroatoms, preferably selected from N, S or O, most preferably selected from N or O, such as

10



R^{10} is a straight chain alkyl of 1 to 6 carbon atoms;

- 15 or a pharmaceutically acceptable salt, ester or prodrug form thereof.

When used herein the term alkyl as a moiety or part of a moiety, e.g. alkoxy, includes straight and branched chain alkyl groups e.g. methyl, ethyl, propyl, i-propyl, n-butyl, s-butyl, i-butyl, t-butyl, pentyl, hexyl and heptyl groups. When used herein the term cycloalkyl includes saturated and unsaturated cyclic groups, e.g. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopropenyl, cyclobutenes, cyclopentenenes, cyclohexenes and cycloheptenes. Saturated cycloalkyl groups are preferred.

- 25 For the compounds defined above and referred to herein, unless otherwise noted, aryl groups include, for example, phenyl and naphthyl which can be substituted, independently, with one or more substituents from the group of hydrogen, halogen, cyano, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, alkoxy of 1 to 6 carbons, CF_3 , or phenyl, the phenyl substituent
30 being further optionally substituted with one or more substituents selected from the

group of hydrogen, halogen, cyano, straight chain alkyl of 1 to 6 carbons, branched chain alkyl of 3 to 7 carbons, or CF₃. The preferred aryl groups herein are phenyl, biphenyl, and naphthyl, substituted or unsubstituted.

5 For the compounds defined above and referred to herein, unless otherwise noted, aroyl groups include, for example, benzoyl and naphthoyl which can be substituted, independently, with one or more substituents from the group of hydrogen, halogen, cyano, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, alkoxy of 1 to 6 carbons, CF₃, or phenyl, the phenyl substituent
10 being further optionally substituted with one or more substituents selected from the group of hydrogen, halogen, cyano, straight chain alkyl of 1 to 6 carbons, branched chain alkyl of 3 to 7 carbons, or CF₃. The preferred aroyl groups herein are benzoyl, (phenyl)benzoyl and naphthoyl, substituted or unsubstituted.

15 For the compounds defined above and referred to herein, unless otherwise noted, heteroaroyl refers to a five or six membered heterocyclic ring directly bonded to a carbonyl radical, and containing one or two heteroatoms selected from nitrogen, oxygen or sulfur, for example thiophene carbonyl or pyridine carbonyl. The heteroaroyl herein can be substituted, independently, with one or more substituents
20 from the group of hydrogen, halogen, cyano, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbons, or CF₃.

 For the compounds defined above and referred to herein, unless otherwise noted, arylalkanoyl refers to groups such as benzyl carbonyl or naphthylmethyl
25 carbonyl containing an alkyl group of 1 to 6 carbon atoms directly bonded to a carbonyl radical, wherein the alkyl group is terminally substituted by an aryl, and the aryl group is as defined hereinbefore.

 For the compounds defined above and referred to herein, unless otherwise
30 noted, aralkyl or arylalkyl refers to groups such as benzyl or naphthylmethyl containing an alkyl residue, preferably a lower alkyl residue of from 1 to 6 carbon atoms, most preferably from 1 to 3 carbon atoms, terminally substituted by an aryl, wherein the aryl group is as defined hereinbefore.

For the compounds defined above and referred to herein, unless otherwise noted, the term halogen is meant to include chlorine, bromine, fluorine and iodine.

The preferred compounds of this invention include:

- 5 [2-Chloro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-methanone;
- [2-Chloro-4-(5-methyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-methanone;
- 10 [2-Bromo-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone;
- 15 (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-(4-fluoro-2-trifluoromethyl-phenyl)-methanone;
- (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[4-(3-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-methanone;
- 20 (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[4-(3-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-methanone 1:1 salt with methanesulfonic acid;
- (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[4-(3-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-methanone 1:1 salt with hydrochloric acid;
- 25 4-(3-Methyl-pyrazol-1-yl)-2-trifluoromethyl-benzoic acid methylester;
- 4-(3-Methyl-pyrazol-1-yl)-2-trifluoromethyl-benzoic acid;
- 30 (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[4-(5-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-methanone;
- (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[2-trifluoromethyl-4-(3-trifluoromethyl-pyrazol-1-yl)-phenyl]-methanone;
- 35

- (5-Methyl-5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[4-(3-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-methanone;
- 5 (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[2-fluoro-4-(3-methyl-pyrazol-1-yl)-phenyl]-methanone;
- (2,4-Difluoro-phenyl)-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone;
- 10 (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[4-fluoro-2-(3-methyl-pyrazol-1-yl)-phenyl]-methanone;
- [2-Chloro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5-methyl-5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone;
- 15 (2-Chloro-4-fluoro-phenyl)-(5-methyl-5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone;
- 20 (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[2-methyl-5-(3-methyl-pyrazol-1-yl)-phenyl]-methanone;
- (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-(5-fluoro-2-methyl-phenyl)-methanone;
- 25 [4-(3-tert-Butyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone;
- [2-Chloro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(11H-5-oxa-4,10-diaza-dibenzo[a,d]cyclohepten-10-yl)-methanone;
- 30 [2-Chloro-4-(3-trifluoromethyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-methanone;
- 35 [2-Chloro-4-(1-methyl-1H-pyrazol-3-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]

]benzodiazepin-10-yl)-methanone;

[2-Chloro-4-(1-methyl-1H-pyrazol-3-yl)-phenyl]-(5-methyl-5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone;

5

6,11-Dihydro-5H-pyrido-[2,3-b][1,5]-benzodiazepine-5-one;

[2-Chloro-4-(5-methyl-1H-[1,2,4]triazol-3-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-methanone;

10

[2-Bromo-4-(3-methyl-pyrazol-1-yl)-phenyl]-(11H-5-oxa-4,10-diazadibenzo[a,d]cyclohepten-10-yl)-methanone;

[4-(3-Methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-(11H-5-oxa-4,10-diazadibenzo[a,d] cyclohepten-10-yl)-methanone;

15

[2-Fluoro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(11H-5-oxa-4,10-diazadibenzo[a,d]cyclohepten-10-yl)-methanone; and

20 [2-Chloro-4-(1-methyl-1H-pyrazol-3-yl)-phenyl]-(11H-5-oxa-4,10-diazadibenzo[a,d]cyclohepten-10-yl)-methanone.

It is understood by those practicing the art that some of the compounds of this invention depending on the definition of R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , X, Y, and Z may contain one or more asymmetric centers and may thus give rise to optical isomers and diastereomers. The present invention includes such optical isomers and diastereomers; as well as the racemic and resolved, enantiomerically pure R and S stereoisomers; as well as other mixtures of the R and S stereoisomers and pharmaceutically acceptable salts thereof, which possess the indicated activity. Optical isomers may be obtained in pure form by standard procedures known to those skilled in the art. It is also understood that this invention encompasses all possible regioisomers, and mixtures thereof which possess the indicated activity. Such regioisomers may be obtained in pure form by standard separation procedures known to those skilled in the art.

35

The pharmaceutically acceptable salts are those derived from such organic and inorganic acids as: citric, lactic, acetic, tartaric, succinic, maleic, malonic, hydrochloric, hydrobromic, phosphoric, nitric, sulfuric, methanesulfonic, and similarly known acceptable acids.

5

Also according to the present invention there is provided a method of treating diseases, conditions or disorders in which vasopressin agonist activity is desired, the method comprising administering to a human or other mammal in need thereof an effective amount of a compound or a pharmaceutical composition of this invention.

10 The present methods of treatment include those for diseases, conditions or disorders which make it desirable to release factor VIII and von Willebrand factor into the circulatory system, release tissue-type plasminogen activator (t-PA) in the blood circulation, or affect the renal conservation of water and urine concentration. Such methods of treatment include, but are not limited to, treatments for diabetes insipidus,
15 nocturnal enuresis, nocturia, urinary incontinence, or bleeding and coagulation disorders in humans or other mammals.

The methods herein include facilitation in humans or other mammals of temporary delay of urination, which may also be described as controlling or treating
20 the inability to temporarily delay urination, whenever desirable. This method is understood to include treatments facilitating the temporary delay of urination which are separate from and not included in the treatment of the conditions known as nocturnal enuresis and nocturia.

25 The present invention accordingly provides a pharmaceutical composition which comprises a compound of this invention in combination or association with a pharmaceutically acceptable carrier. In particular, the present invention provides a pharmaceutical composition which comprises an effective amount of a compound of this invention and a pharmaceutically acceptable carrier or excipient.

30

The compositions are preferably adapted for oral administration. However, they may be adapted for other modes of administration, for example, parenteral administration for patients suffering from heart failure.

In order to obtain consistency of administration, it is preferred that a composition of the invention is in the form of a unit dose. Suitable unit dose forms include tablets, capsules and powders in sachets or vials. Such unit dose forms may contain from 0.1 to 1000 mg of a compound of the invention and preferably from 2 to 50 mg. Still further preferred unit dosage forms contain 5 to 25 mg of a compound of the present invention. The compounds of the present invention can be administered orally at a dose range of about 0.01 to 100 mg/kg or preferably at a dose range of 0.1 to 10 mg/kg. Such compositions may be administered from 1 to 6 times a day, more usually from 1 to 4 times a day. The compositions of the invention may be formulated with conventional excipients, such as a filler, a disintegrating agent, a binder, a lubricant, a flavoring agent and the like. They are formulated in conventional manner, for example, in a manner similar to that used for known antihypertensive agents, diuretics and β -blocking agents.


Also according to the present invention there are provided processes for producing the compounds of the present invention.

PROCESS OF THE INVENTION

The compounds of the present invention may be prepared according to one of the general processes outlined below. The compounds of general formula (I) wherein W is oxygen or NR⁶ and R⁶ is hydrogen, can be conveniently prepared as shown in Scheme I.

-17-

SCHEME I




Thus, a pyridobenzodiazepine (or benzoxazepine) of formula (1) is treated with an appropriately substituted haloaroyl(heteroaroyl) halide, preferably a fluoroaroyl or a fluoro(chloro)heteroaroyl chloride of formula (2, J= COCl), in the presence of an inorganic base such as potassium carbonate in a polar, aprotic solvent such as dimethylformamide; or an organic base such as 4-dimethylaminopyridine in an aprotic solvent such as dichloromethane or tetrahydrofuran at temperatures ranging from -40°C to 50°C to yield the intermediate acylated derivative (3).

10

Alternatively, the acylating species can be a mixed anhydride of the corresponding carboxylic acid, such as that prepared treating said acid with 2,4,6-trichlorobenzoyl chloride in an aprotic organic solvent such as dichloromethane according to the procedure of Inanaga et al., *Bull. Chem. Soc. Jpn.*, **52**, 1989 (1979).

Treatment of said mixed anhydride of general formula (2) with the pyridobenzodiazepine (or benzoxazepine) of formula (1) in a solvent such as dichloromethane and in the presence of an organic base such as 4-dimethylaminopyridine at temperatures ranging from 0°C to the reflux temperature of the solvent, yields the intermediate acylated derivative (3) of Scheme I.

A compound of formula (3) is then treated with the sodium (potassium or lithium) salt of an appropriately substituted heterocycle of formula (4, wherein R¹ is selected from the a, b, c, d, l, n, or o group of heterocycles defined above) in an aprotic organic solvent such as dimethylformamide (or tetrahydrofuran) at temperatures ranging from ambient to the reflux temperature of the solvent, to yield a compound of general formula (I), wherein W is oxygen or NH, and A, B, X, Y, Z, R², R³, and R⁵ are as defined above, R⁶ is hydrogen, and R¹ is an heterocyclic moiety selected from the a, b, c, d, l, n or o group of heterocycles defined above and illustrated below.



The condensation of the intermediate of formula (3) with the intermediate salt of formula (4) leads to a variable ratio of regioisomers of general formula (I) which are separated by means of chromatography and/or crystallization.

The preferred substituted fluoroaroyl and fluoro(chloro)heteroaroyl chlorides of formula (2) of Scheme I are either available commercially, or are known in the art, or can be readily prepared by procedures analogous to those in the literature for the known compounds.

5


The sodium (potassium or lithium) salts of the heterocycle of formula (4, wherein R¹ is selected from the a, b, c, d, l, n or o group of heterocycles defined above) of Scheme I are prepared by treatment of said heterocycle with a strong base such as sodium, potassium or lithium hydride or a metal alkoxide at temperatures ranging from -40°C to ambient in an aprotic organic solvent such as dimethylformamide or tetrahydrofuran.

10

Alternatively, the compounds of general formula (I) described in Scheme I can be prepared according to the process outlined in Scheme II.

15

SCHEME II



Thus, an appropriately substituted fluoroaryl or fluoro(chloro)heteroaryl carboxylic acid of formula (5) is esterified using methods known in the art, such as treatment with oxalyl chloride or thionyl chloride in an alcohol solvent such as methanol, in the presence of a catalytic amount of dimethylformamide; or by condensation with an alcohol such as methanol, in the presence of an acid catalyst such as para-toluenesulfonic acid at temperatures ranging from ambient to reflux temperature of the solvent.

5 The resulting ester of formula (6) is reacted with the sodium (potassium or lithium) salt of an appropriately substituted heterocycle of formula (4, wherein R¹ is selected from the a, b, c, d, l, n or o group of heterocycles defined above) in a polar aprotic organic solvent such as dimethylformamide at temperatures ranging from ambient to reflux temperature of the solvent, to yield an intermediate ester of general formula (7). The condensation of (6) with (4) leads to a variable ratio of regioisomers of general formula (7) which are separated by means of chromatography and/or crystallization.

10 Subsequent hydrolysis of the intermediate ester of formula (7) with an aqueous base such as sodium hydroxide in methanol (or lithium hydroxide in tetrahydrofuran) affords the carboxylic acid of general formula (8).


15 The intermediate carboxylic acid (8) is then converted into an acylating agent, preferably an acid chloride or a mixed anhydride of general formula (9) using any of the procedures described hereinbefore.

20 Subsequent condensation of the pyridobenzodiazepine (or benzoxazepine) of formula (1) with the intermediate acylating agent of formula (9) according to any of the procedures described hereinbefore, yields the desired compounds of formula (I) of Scheme I, wherein R¹ is selected from the a, b, c, d, l, n or o group of heterocycles defined above.

25 The appropriately substituted fluoroaryl or fluoro(chloro)heteroaryl carboxylic acids of formula (5) of Scheme II are either available commercially, or are known in the art, or can be readily prepared by procedures analogous to those in the literature for known compounds.

30 Alternatively, the substituted carboxylic acids of formula (8) of Scheme II can be prepared according to the process outlined in Scheme III.

SCHEME III



Thus, a fluoroaryl or fluoro(chloro)heteroaryl nitrile of formula (10) is reacted with the sodium (potassium or lithium) salt of an appropriately substituted heterocycle of formula (4, wherein R¹ is selected from the a, b, c, d, l, n or o group of heterocycles defined above) in a polar aprotic organic solvent such as dimethylformamide at temperatures ranging from ambient to the reflux temperature of the solvent, to yield an intermediate of general formula (11). Condensation of (10) with the intermediate (4) leads to a variable ratio of regioisomers of general formula (11) which are separated by means of chromatography and/or crystallization. Hydrolysis of the intermediate nitrile of formula (11, wherein Y is not CF₃) is preferentially carried out with an inorganic acid such as dilute sulfuric acid, at temperatures ranging from ambient to 60°C.

Alternatively, hydrolysis of the nitrile (11) can be carried out by heating in the presence of a strong alkaline base such as sodium hydroxide in an alcohol solvent such as ethanol, with or without a phase transfer catalyst such as benzyldimethyltetradecyl ammonium chloride.


The resulting carboxylic acid of formula (8) is then converted into the desired compounds of formula (I) of Scheme I (wherein R¹ is selected from the a, b, c, d, l, n and o group of heterocycles defined above) by procedures analogous to those described hereinbefore.

The appropriately substituted fluoroaryl or fluoro(chloro)heteroaryl nitriles of formula (10) of Scheme III are either available commercially, or are known in the art,

or can be readily prepared by procedures analogous to those in the literature for known compounds.

Alternatively, the intermediate substituted carboxylic acids of formula (8, wherein R^1 is not **b** or **d**) of Scheme II can be prepared according to the process described in Scheme IV by sequential treatment of a nitrile of formula (11, wherein A and B are carbon, and R^1 is an heterocyclic moiety selected from the **a**, **c**, **e**, **l**, **n** and **o**, but not **b** or **d** group of heterocycles defined above) with basic hydrogen peroxide in dimethylsulfoxide essentially according to the procedure of Katritzky et al., *Synthesis*, 949 (1989); followed by hydrolysis of the resulting amides of formula (12) preferably by treatment with dilute sulfuric acid and sodium nitrite according to the procedure of Hales et al, *Tetrahedron*, 51, 7403 (1995).


SCHEME IV



15


A preferred process for the preparation of the intermediate substituted carboxylic acids of formula (8) of Scheme II wherein R^1 is an heterocyclic moiety selected from the **a** group of R^1 heterocycles defined above, is outlined in Scheme V.

SCHEME V



Thus, diazotization of an appropriately substituted aniline of general formula (13) followed by reduction of the resulting diazonium salt of formula (14) with tin (II) chloride in concentrated hydrochloric acid according to the procedure of Street et al., *J. Med. Chem.*, **36**, 1529 (1993) provides the intermediate hydrazine hydrochloride salt of formula (15). Subsequent condensation of (15) with an aldehyde derivative of formula (35, wherein R^2 and R^5 are as defined above, R^3 is hydrogen, and P is a dialkylacetal) such as acetylacetaldehyde dimethyl acetal, or a ketone (or a ketone derivative) of formula (35, wherein R^2 and R^5 are as defined above, R^3 is not hydrogen, and P is O or a ketal) in a solvent such as aqueous methanol at temperatures


- 5 ranging from ambient to 100°C provides after crystallization, the desired intermediate ester of formula (7, wherein R¹ is selected from the a group of heterocycles defined above), which is then converted to the compound of formula (I, wherein R¹ is selected from the a group of heterocycles defined above and illustrated below) as outlined in Scheme II above



- 10 The compounds of general formula (I) wherein W is NR⁶ and R⁶ is other than hydrogen, and R¹ is selected from the a, b, c, d, l, n, or o group of heterocycles defined above, can be prepared by alkylation or acylation of the intermediate of formula (3, W=NH) of Scheme I, as outlined in Scheme VI.

-26-


SCHEME VI

**(3 W= NH)**

Thus, the intermediate compound (3, W= NH) of Scheme I is alkylated by
 5 treatment with a base such as sodium (or potassium) hydride and an alkylating agent
 such as an alkyl halide, preferably an alkyl chloride (bromide or iodide), in an aprotic
 solvent such as dimethylformamide or tetrahydrofuran at temperatures ranging from
 0°C to 80°C to yield compounds of formula (16, wherein A, B, X, Y, Z, and R⁶ are as
 defined above).

Alternatively, a compound of formula (3, W =NH) of Scheme I is acylated by treatment with a carboxylic acid halide or a carboxylic acid anhydride in the presence of an amine base such as pyridine or triethylamine in an aprotic solvent such as dichloromethane or with no addition of solvent when pyridine is used as the base, at temperatures ranging from -40°C to ambient, to yield compounds of structure (16 wherein A, B, X, Y, Z, and R⁶ are as defined above).


Subsequent treatment of the compounds of formula (16, wherein R⁶ is either an alkyl or acyl moiety) with the sodium (potassium or lithium) salt of an appropriately substituted heterocycle of formula (4, wherein R¹ is selected from the a, b, c, d, l, n or o group of heterocycles defined above) in a polar aprotic organic solvent such as dimethylformamide at temperatures ranging from ambient to reflux temperature of the solvent, yields compounds of general formula (I), wherein W is NR⁶, and R⁶ is an alkyl or acyl residue; A, B, X, Y, Z, R², R³, and R⁵ are as defined above, and R¹ is an heterocyclic moiety selected from the group consisting of a, b, c, d, l, n or o, defined above and illustrated below.



The alkylating and acylating agent(s) of Scheme VI are either available commercially, or are known in the art, or can be readily prepared by procedures analogous to those in the literature for known compounds.

Alternatively, a compound of general formula (I) of Scheme VI, can be prepared by a one-pot process outlined in Scheme VII.


SCHEME VII




Thus, a fluoroaryl or fluoro(chloro)heteroaryl intermediate of formula (3) of Scheme I (wherein W is NH) is reacted with the sodium (potassium or lithium) salt of an appropriately substituted heterocycle of formula (4, wherein R^1 is selected from the a, b, c, d, l, n or o group of heterocycles defined above) in a polar aprotic organic solvent such as dimethylformamide at temperatures ranging from ambient to the reflux temperature of the solvent, generating an intermediate of general formula (I, wherein W is NH) which without prior isolation, is further alkylated (or acylated) using the reaction conditions described hereinbefore to provide the desired compound of formula (I, wherein W is NR^6 , and R^6 is an alkyl or acyl residue). This process leads also to a variable ratio of regioisomers of general formula (I), wherein W is NR^6 , and R^6 is an alkyl or acyl residue; A, B, X, Y, Z, R^2 , R^3 , R^4 and R^5 are as defined above, and R^1 is a heterocyclic moiety defined as in Scheme VI. The regioisomers of formula (I) can be separated by means of chromatography and/or crystallization.

The compounds of general formula (I) of Scheme I wherein A and B are carbon, R^2 is hydrogen, and R^1 is a heterocyclic moiety selected from the g group of heterocycles described above, may be prepared according to the general process outlined in Scheme VIII.

SCHEME VIII



-30-



I (W=O, NH; A,B= carbon; R¹= g)

Thus, an appropriately substituted haloaryl carboxylic acid ester, preferably a bromo (or iodo)phenyl methylester of formula (17, wherein A and B are carbon) is coupled with a dialkylamino propyne, in the presence of a catalyst such as bis(triphenylphosphine) palladium (II) chloride and copper (I) iodide in an organic base such as triethylamine as the solvent, and at temperatures ranging from ambient to 80°C essentially according to the procedures of Alami et al., *Tetrahedron Lett.*, **34**, 6403 (1993), and of Sanogashira et al., *Tetrahedron Lett.*, 4467 (1975), to provide the substituted acetylene intermediate of general formula (18).

10

The intermediate (18) is subsequently converted into its N-oxide by treatment with an oxidizing agent using any of a number of standard oxidative procedures (see Albini, *Synthesis*, 263, (1993)) or with a dioxirane reagent (see Murray, *Chem. Rev.*, 1187 (1989)) in an aprotic organic solvent such as dichloromethane at temperatures below ambient. The intermediate N-oxide is not isolated but is rearranged *in situ* to an enone of general formula (19) by heating in a hydroxylic solvent such as methanol or by using any of the procedures of Dushin et al. This procedure provides a novel synthesis of enaminone compounds from propargylic amines (or their N-oxides) in hydroxylic solvents, which influence the ultimate outcome of the reaction. This new method of enaminone synthesis provides a convenient alternative to existing methods and further extends the range of starting materials that can be converted into enaminone products.

15

20


Although the precise mechanism by which a propargylic amine N-oxide is converted into an enaminone product has not been rigorously determined, it likely resembles two known processes; the thermal [2,3]-sigmatropic rearrangement of

25

propargylic amine N-oxides (Craig, et al. *Tetrahedron Lett.*, 4025, (1979); Hallstrom, et al., *Tetrahedron Lett.*, 667, (1980); Khuthier, A-H, et al., *J. Chem. Soc. Chem. Commun.*, 9, (1979) and the conversion of certain isoxazoles into enaminones (Liguori, et al., *Tetrahedron*, 44, 1255 (1988).


5 Treatment of (19) with a substituted hydrazine (36) in acetic acid at temperatures ranging from ambient to reflux leads to a mixture of regioisomeric compounds of general formulas (20) and (21) in a variable ratio. The major isomer of formula (20, wherein R^2 is H) is separated by means of chromatography and/or
10 crystallization and is subsequently hydrolyzed to the desired carboxylic acid of formula (22).

The intermediate (22) is then converted into an acylating species, preferably an acid chloride (bromide or iodide) or a mixed anhydride of formula (23) by
15 procedures analogous to those described hereinbefore. The acylating agent (23) is then used to acylate a pyridobenzodiazepine (or benzoxazepine) of formula (1) by any of the procedures described hereinbefore to yield the desired compound of formula (I), wherein W is O or NH, A and B are carbon, X, Y, Z, and R^4 are as defined above, R^2 is hydrogen, and R^1 is an heterocyclic moiety selected from the g group of
20 heterocycles defined above and illustrated below.




Likewise, treatment of (19) with an unsubstituted hydrazine (36, wherein R^4 is H) in acetic acid at temperatures ranging from ambient to the reflux temperature of the solvent yields the intermediate pyrazole ester of formula (24, R^2 and R^4 are H) as
25 shown in Scheme IX. In this case the heterocyclic nitrogen of (24) can be alkylated or acylated to provide intermediates which can be converted to compounds of formula (I) wherein W is O or NH, A and B are carbon, X, Y, Z, and R^4 are as defined above, R^2 is hydrogen, and R^1 is an heterocyclic moiety selected from the f group of heterocycles defined above.

SCHEME IX




Thus, the intermediate ester of formula (24, wherein R² is H, A and B are carbon) is alkylated by treatment with a base such as sodium or potassium hydride

and an alkylating agent such as an alkyl halide, preferably an alkyl chloride (bromide or iodide) in an aprotic solvent such as dimethylformamide or tetrahydrofuran at temperatures ranging from 0°C to 80°C to yield a mixture of regioisomers of formulas (20) and (21) in a variable ratio. The major regioisomer of formula (21) is
5 separated by chromatography and/or crystallization and is subsequently hydrolyzed to the desired carboxylic acid of formula (25), which is then converted to an acylating agent, preferably an acid chloride or a mixed anhydride by procedures analogous to those described hereinbefore. The acylating species of formula (26) is then used to acylate a pyridobenzodiazepine (or benzoxazepine) of formula (1) to yield the
10 desired compound of formula (I), wherein W is O or NH, A and B are carbon, X, Y, Z, and R⁴ are as defined above, R² is hydrogen, and R¹ is a heterocyclic moiety selected from the f group of heterocycles defined above and illustrated below.




Compounds of general formula (I) wherein W= is O or NH, A and B are
15 carbon, and R¹ is a heterocyclic moiety selected from the h group of heterocycles defined above, can be prepared as outlined in Scheme X.

SCHEME X



- An appropriately substituted malondialdehyde of formula (27) is treated first with a hydrazine in acetic acid at temperatures ranging from ambient to the reflux temperature of the solvent and the intermediate pyrazole is oxidized preferably with potassium permanganate in a basic aqueous solution at temperatures ranging from ambient to the reflux temperature of the solvent to yield a carboxylic acid intermediate of formula (28). The acid (28) is converted into an acylating agent, preferably an acid chloride (bromide or iodide) or a mixed anhydride by procedures analogous to those described hereinbefore. The acylating agent of formula (29) is finally reacted with a pyridobenzodiazepine (or benzoxazepine) of formula (1) to yield compounds of general formula (I) wherein W is O or NH, A and B are carbon, X, Y, Z, and R⁴ are as defined above, and R¹ is an heterocyclic moiety selected from the h group of heterocycles defined above and illustrated below.


-35-



The preferred malondialdehydes of formula (27) and the hydrazines (36) of Scheme X are either available commercially, or are known in the art, or can be readily prepared by procedures analogous to those in the literature for known compounds, such as those of Knorr et al., *J. Org. Chem.*, 49, 1288 (1984) and Coppola et al., *J. Het. Chem.*, 51 (1974).

An alternative preparation of the intermediate carboxylic acids of formula (28) of Scheme X wherein Y is as defined above and R⁴ is other than hydrogen, is outlined in Scheme XI.

SCHEME XI



An organotin reagent of formula (31) is reacted in a Stille coupling reaction with an appropriately substituted aryl(heteroaryl) halide, preferably a bromide (or iodide) of formula (34) in the presence of a catalyst such as tetrakis(triphenylphosphine) palladium (0) and copper (I) iodide in an organic aprotic solvent such as dimethylformamide at temperatures ranging from ambient to 150°C, essentially according to procedures analogous to those of Farina et al., *J. Org. Chem.*, 59, 5905 (1994). Basic hydrolysis of the resulting ester of formula (32) with sodium hydroxide in aqueous alcohol or lithium hydroxide in aqueous tetrahydrofuran at temperatures ranging from ambient to the reflux temperature of the solvent yields the desired carboxylic acids of formula (28) of Scheme X.

In turn, the organotin reagents of formula (31) of Scheme XI wherein R is preferably an alkyl group, are conveniently prepared by metallation of a 4-bromo-N-alkylpyrazole of formula (30) with a trialkyltin halide, preferably a tributyltin chloride (or bromide) in the presence of a metallating agent such as n-butyl lithium in an aprotic organic solvent such as diethyl ether at temperatures ranging from -40°C to ambient according to procedures analogous to those found in Martina et al., *Synthesis*, 8, 613 (1991).


The preferred N-alkyl substituted pyrazoles of formula (30) of Scheme XI are conveniently prepared from a 4-bromopyrazole by alkylation with an alkyl halide, preferably an alkyl chloride (bromide or iodide) in the presence of a base such as sodium (or potassium) hydride in an aprotic organic solvent such as dimethylformamide or tetrahydrofuran at temperatures ranging from 0°C to 80°C. Alternatively, alkylation of the 4-bromopyrazole can be carried out with an alkylating agent mentioned above, and a strong alkaline base such as sodium hydroxide in the presence of a phase transfer catalyst, such as benzyldimethyltetradecyl ammonium chloride or benzyltrimethylammonium chloride (see Jones, *Aldrichimica Acta*, 9, 35 (1976)).

The preferred aryl (heteroaryl) iodides of formula (34) of Scheme XI are conveniently prepared by diazotization of the corresponding substituted anilines of formula (33) followed by reaction of the corresponding diazonium salt with iodine and potassium iodide in aqueous acidic medium essentially according to the

procedures of Street et al., *J. Med. Chem.*, **36**, 1529 (1993) and of Coffen et al., *J. Org. Chem.*, **49**, 296 (1984).

The compounds of general formula (I) may be prepared also according to one
5 of the general processes outlined below.

As shown in Scheme XII, a pyridobenzodiazepine (or benzoxazepine) of
formula (1) is treated with an appropriately substituted acetylaroyl (heteroaroyl)halide
preferably an acetylaroyl (heteroaroyl)chloride of formula (37, J=COCl) according to
10 any of the procedures described hereinbefore, to yield the acylated derivative of
formula (38). Treatment of (38) with a dialkylamide dialkylacetal such as a
dimethylamide dimethyl acetal of formula (39, where alkyl is CH₃), in an aprotic
organic solvent such as dichloromethane at temperatures ranging from 0°C to the
reflux temperature of the solvent according to the procedure of Lin et al., *J. Het.*
15 *Chem.*, 345 (1977) yields the enone of formula (40). Treatment of (40) with
hydroxylamine or a substituted hydrazine of formula (36) in acetic acid at
temperatures ranging from ambient to the reflux temperature of the solvent provides
the target compounds of formula (I) wherein W is O or NH, A, B, X, Y, Z, R² and
R⁴ are as defined above, and R¹ is an heterocyclic moiety selected from the f, g, or j
20 group of heterocycles defined above and illustrated below.




The preferred substituted acetylaroyl (heteroaroyl)chlorides of formula (37) of
25 Scheme XII are conveniently prepared by treating the corresponding carboxylic acids
with thionyl chloride at temperatures ranging from ambient to the reflux temperature
of the solvent, or with oxalyl chloride in an aprotic solvent such as dichloromethane
or tetrahydrofuran in the presence of a catalytic amount of dimethylformamide at
temperatures ranging from 0°C to 40°C.

30

The preferred dialkylamide dialkylacetals of formula (39) of Scheme XII are
either available commercially, or are known in the literature, or can be conveniently

prepared according to procedures analogous to those in the literature for the known compounds (see Kantlehner, *Chem. Ber.*, **105**, 1340 (1972)).


SCHEME XII



5

An alternate process for the preparation of intermediates of formula (38) of Scheme XII is illustrated in the following Scheme XIII.

SCHEME XIII



Thus, a pyridobenzodiazepine (or benzoxazepine) of formula (1) is treated with an appropriately substituted bromoaryl(heteroaryl) halide, preferably a bromoaryl(heteroaryl) chloride of formula (42) according to any of the procedures described hereinbefore, to yield the acylated intermediate of formula (43). The intermediate (43) is subsequently coupled with a monosubstituted acetylene of formula (53, wherein R^9 is preferably trimethylsilyl or lower alkyl of 1 to 6 carbon atoms) in the presence of pyridine and a catalyst such as bis(triphenylphosphine) palladium (II) chloride and copper (I) iodide in an organic base such as triethylamine as the solvent, in a sealed pressure tube at temperatures ranging from ambient to 100°C essentially according to the procedure of Martinez et al., *J. Med. Chem.*, **52**, 3491 (1987). The resulting acetylene intermediate of formula (44) is then hydrated by treatment with 1% sulfuric acid in an aprotic organic solvent such as tetrahydrofuran saturated with mercury (II) sulfate at ambient temperature essentially according to the procedure of Reed et al., *J. Org. Chem.*, **52**, 3491 (1987) to provide the desired acyl compound of formula (38) wherein W is O or NH, and A, B, X, Y, and Z are as defined above, and R^9 is hydrogen or lower alkyl of 1 to 6 carbon atoms.

Alternatively, compound (44) where R⁹ is trimethylsilyl is treated with n-tetrabutylammonium fluoride in an ether solvent such as tetrahydrofuran, to afford compound (44) where R⁹ is hydrogen.


5 The preferred acylating agents of formula (42) of Scheme XIII are conveniently prepared by treating an appropriately substituted aryl(heteroaryl) carboxylic acid of formula (41) with thionyl chloride at temperatures ranging from ambient to the reflux temperature of the solvent, or with oxalyl chloride in an aprotic solvent such as dichloromethane or tetrahydrofuran in the presence of a catalytic
10 amount of dimethylformamide at temperatures ranging from 0°C to 40°C.

The acetylene intermediates (53) of Scheme XIII are either available commercially, or are known in the art, or can be readily prepared by procedures analogous to those in the literature for the known compounds.

15

As shown in Scheme XIV, the intermediate acetyl compounds (38) of Scheme XII can be prepared also by the Stille coupling of a bromo aryl (heteroaryl) compound of formula (43) of Scheme XIII with an (α -alkoxyvinyl)trialkyltin preferably an (α -ethoxyvinyl)tributyltin of formula (45), in the presence of a catalytic
20 amount of bis(triphenylphosphine) palladium (II) chloride in an aprotic organic solvent such as toluene at temperatures ranging from ambient to the reflux temperature of the solvent, essentially according to the procedure of Kosugi et al., *Bull. Chem. Soc. Jpn.*, **60**, 767 (1987).

Scheme XIV



25


The preparation of the acetyl compound (38) can also be accomplished via the palladium-catalyzed arylation of a vinyl alkylether such as vinyl butylether, with the

aryl halide intermediate of formula (43) according to the procedure of Cabri et al., *Tetrahedron Lett.*, **32**, 1753 (1991).

5 The (α -alkoxyvinyl)trialkyltin intermediates (45) of Scheme XIV are either available commercially, or are known in the art, or can be readily prepared by procedures analogous to those in the literature for the known compounds.




Compounds in which R¹ contains three heteroatoms are prepared according to Scheme XV.

SCHEME XV




Thus, a pyridobenzodiazepine (or benzoxazepine) of formula (1) is treated with an appropriately substituted cyanoaroyl (heteroaroyl) halide, preferably a cyanoaroyl(heteroaroyl) chloride of formula (46) according to any of the procedures described hereinbefore, to yield an intermediate nitrile of formula (54) which in turn, is converted to an amide intermediate of general formula (47) by treatment with an

inorganic acid such as sulfuric acid at temperatures ranging from ambient to 50°C. Treatment of the amide (47) with a dialkylamide dialkylacetal such as a dimethylamide dimethylacetal of formula (39, wherein alkyl is CH₃) in an aprotic organic solvent such as dichloromethane or tetrahydrofuran at temperatures ranging from 0°C to 80°C yields the intermediate of formula (48). Treatment of (48) with hydroxylamine or a hydrazine of formula (36) in acetic acid at temperatures ranging from ambient to reflux yields the desired target compounds of formula (I) wherein W is O or NH, A, B, X, Y, Z, R² and R⁴ are as defined above, and R¹ is an heterocyclic moiety selected from the e, i or k group of heterocycles defined above and illustrated below.




Another preferred process for the preparation of the intermediate amide of formula (47, where A, B are carbon) of Scheme XV is outlined in Scheme XVI. An appropriately substituted aryl nitrile of formula (49, A and B are carbon) is hydrated with basic hydrogen peroxide in dimethylsulfoxide essentially according to the procedure of Katritzky et al., *Synthesis*, 949 (1989), to provide the intermediate amide (50). Subsequent hydrolysis of the ester moiety gives the carboxylic acid intermediate (51) which is then converted into the acylating species of formula (52) by using any of the procedures described hereinbefore. Treatment of a pyridobenzazepine (or benzoxazepine) of formula (1) with (52) using any of the procedures described hereinbefore provides the desired intermediate amide (47, where A and B are carbon).

SCHEME XVI



Another preferred process to prepare compounds of general formula (I) of Scheme XV where W is O or NH, R¹ is a heterocyclic moiety selected from the e or i group of heterocycles defined above, and R⁴ is not hydrogen, is shown in Scheme XVII.


SCHEME XVII



Thus, an appropriately substituted amide of formula (50) of Scheme XVI is treated with a dialkylamide dialkylacetal such as a dimethylamide dimethylacetal of formula (39, where alkyl is CH₃) at temperatures ranging from 0°C to 100°C to provide the intermediate of formula (55). Treatment of (55) with a substituted hydrazine of formula (36) in acetic acid at temperatures ranging from ambient to reflux, yields the desired intermediate triazole ester of formula (56). The ester (56) is subsequently

- hydrolyzed to the carboxylic acid of formula (57, wherein R^1 is an heterocyclic moiety selected from the e or i group of heterocycles defined above, and R^4 is not hydrogen) which is then converted to an acylating agent, preferably an acid chloride or a mixed anhydride of formula (58) by procedures analogous to those described hereinbefore. The acylating species (58) is used to acylate a pyridobenzodiazepine (or benzoxazepine) of formula (1) to yield the desired compound of formula (I) wherein W is O or NH, A, B, X, Y, Z and R^2 are as defined above, R^1 is an heterocyclic moiety selected from the e and i group of heterocycles defined above and illustrated below, and R^4 is not hydrogen.


10




- Alternatively, the compounds of general formula (I) of Scheme XV wherein W is O or NH, R^1 is an heterocyclic moiety selected from the e or i group of heterocycles defined above, and R^4 is not hydrogen can be prepared as illustrated in Scheme XVIII.

15

SCHEME XVIII




SCHEME XVIII



Treatment of the intermediate ester of formula (55) of Scheme XVII with an unsubstituted hydrazine (36, where R^4 is H) in acetic acid at temperatures ranging from ambient to the reflux temperature, yields the intermediate triazole ester of formula (59). In this case the heterocyclic nitrogen can be alkylated or acylated by procedures analogous to those described hereinbefore, to yield the substituted triazole

ester of formula (60). The ester (60) is subsequently hydrolyzed to the carboxylic acid of formula (61) which is then converted into an acylating species, preferably an acid chloride or mixed anhydride of formula (62), by procedures analogous to those described hereinbefore. The acylating agent (62) is used to acylate a
 5 pyridobenzodiazepine (or benzoxazepine) of formula (1) to yield the desired compound of formula (I) wherein W is O or NH, X, Y, Z, A, B and R² are as defined above, R¹ is a heterocyclic moiety selected from the e or i group of heterocycles defined above, and R⁴ is not hydrogen.

10 Alternatively, a compound of general formula (I) of Scheme XV wherein W is O or NH, A, B, X, Y, Z and R² are as defined above, R¹ is an heterocyclic moiety selected from the e and i group of heterocycles defined above and illustrated below, and R⁴ is hydrogen, can be conveniently prepared from a compound of formula (I)
 15 of Scheme XVIII wherein R⁴ is an optionally substituted aralkyl group, preferably a p-methoxybenzyl group by using a number of procedures which include hydrogenolysis or treatment with a strong acid such as trifluoroacetic acid at temperatures ranging from 0°C to reflux temperature, essentially according to the procedure of Buckle et al., *J. Chem. Soc. Perkin Trans. 1*, 627 (1982).




20 The preferred process to prepare compounds of general formula (I) in which R¹ contains four heteroatoms, W is O or NH, and R⁴ is hydrogen is outlined in Scheme XIX.


25

-50-

SCHEME XIX



5 Treatment of the nitrile intermediate of formula (54) of Scheme XV with sodium azide and ammonium chloride in an aprotic organic solvent such as dimethylformamide at temperatures ranging from ambient to the reflux temperature of the solvent, yields the desired compound of formula (I) wherein W is O or NH, A, B, X, Y, and Z are as defined above, R¹ is a heterocyclic moiety selected from the m group of heterocycles defined above and illustrated below, and R⁴ is hydrogen.




10

m (R⁴ = H)

15 The compounds of general formula (I) wherein R¹ is a heterocyclic moiety selected from the group e, f, g, h, i, j, or k, and W is NR⁶ and R⁶ is other than hydrogen, can be preferably prepared by alkylation or acylation of a compound of formula (I, W is NH) of Schemes VIII, IX, X, XII, and XV, as outlined in Scheme XX.

SCHEME XX




Thus, the compounds of formula (I, W is NH) of Schemes VIII, IX, X, XII, and XV are alkylated by treatment with a base such as sodium (or potassium) hydride and an alkylating agent such as an alkyl halide, preferably an alkyl chloride (bromide or iodide) in an aprotic solvent such as dimethylformamide or tetrahydrofuran at temperatures ranging from 0°C to 80°C to yield compounds of formula (I) wherein W is NR⁶ and R⁶ is alkyl, and A, B and R¹ are as defined in Scheme XX and illustrated below.

10


Alternatively, the compounds of formula (I, W is NH) of Schemes VIII, IX, X, XII, and XV are acylated by treatment with a carboxylic acid halide or a



- carboxylic acid anhydride in the presence of an amine base such as pyridine or a trialkylamine such as triethylamine in an aprotic solvent such as dichloromethane or with no addition of solvent when pyridine is used as the base, at temperatures ranging from -40°C to ambient, to yield compounds of formula (I) wherein W is NR⁶ and R⁶ is acyl, and A, B and R¹ are as defined in Scheme XX and illustrated below.
- 5



- The compounds of general formula (I) wherein R¹ is an heterocyclic moiety selected from the group e, i, k, or m, and W is NR⁶ and R⁶ is other than hydrogen, can be conveniently prepared as shown in Scheme XXI.
- 10

SCHEME XXI






Thus a nitrile of formula (54, W is NH) of Scheme XV can be alkylated by treatment with a base such as sodium (or potassium) hydride and an alkylating agent such as an alkyl halide, preferably an alkyl chloride (bromide or iodide) in an aprotic solvent such as dimethylformamide or tetrahydrofuran at temperatures ranging from 0°C to 80°C to yield an alkylated nitrile of formula (63, W is NR⁶ and R⁶ is alkyl).

Conversely (54) can be acylated by treatment with a carboxylic acid halide or a carboxylic acid anhydride in the presence of an amine base such as pyridine or a trialkylamine such as triethylamine in an aprotic solvent such as dichloromethane or with no addition of solvent when pyridine is used as the base, at temperatures ranging from -40°C to ambient, to yield compounds of formula (63, W is NR⁶ and R⁶ is acyl).

In a likely fashion using procedures analogous to those used for the nitrile (54, W= NH) above, an amide of formula (47, W is NH) of Scheme XV can be alkylated or acylated to yield alkylated or acylated intermediates of general formula (64, W= NR⁶ and R⁶= alkyl or acyl residue).

The intermediate nitrile (63) and amide (64) of Scheme XXI can then be converted respectively to either the compound of formula (I), wherein W is NR⁶ and R⁶= alkyl or acyl, and R¹ is selected from the group m of heterocycles defined in Scheme XIX, or to the compound of formula (I) wherein W= NR⁶ and R⁶ is alkyl or acyl and R¹ is selected from the group e, i or k of heterocycles defined in Scheme XV

and illustrated below, by using procedures identical to those outlined in those same Schemes.



5

The subject compounds of the present invention were tested for biological activity according to the following procedures.

Vasopressin V₂ Agonist Effects of Test Compounds in Normal Conscious Water-

Loaded Rats:

Male or female normotensive Sprague-Dawley rats (Charles River Laboratories, Inc., Kingston, NY) of 350-500 g body weight were supplied with standard rodent diet (Purina Rodent Lab. Chow 5001) and water ad libitum. On the day of test, rats were placed individually into metabolic cages equipped with devices to separate the feces from the urine and containers for collection of urine. A test compound or a reference agent was given at an oral dose of 10 mg/kg in a volume of 10 ml/kg. The vehicle used was 20% dimethylsulfoxide (DMSO) in 2.5% preboiled corn starch. Thirty minutes after dosing the test compound, rats were gavaged with water at 30 ml/kg into the stomach using a feeding needle. During the test, rats were not provided with water or food. Urine was collected for four hours after dosing of the test compound. At the end of four hours, urine volume was measured. Urinary osmolality was determined using a Fiske One-Ten Osmometer (Fiske Associates, Norwood, MA, 02062) or an Advanced CRYOMATIC Osmometer, Model 3C2 (Advanced Instruments, Norwood, MA). Determinations of Na⁺, K⁺ and Cl⁻ ion were carried out using ion specific electrodes in a Beckman SYNCHRON EL-ISE Electrolyte System analyzer. The urinary osmolality should increase proportionally. In the screening test, two rats were used for each compound. If the difference in the urine volume of the two rats was greater than 50%, a third rat was used. The results of this study are shown in Table I, below.

Table 1

Example	Urine Volume (% decrease) ^a	Changes in Urinary Osmolality ^b	Rat Type ^c
1	80	272	CD
2	61	288	CD
3	85	346	CD
4	87	339	CD
6	25	103	CD
7	69	283	CD
8	72	298	CD
9	59	372	CD
10	63	276	CD
11	71	375	CD
12	-2	101	CD
13	6	106	CD
14	83 (1 mg/kg)	321	CD
15	98 (1 mg/kg) starch only	1363	CD
16	41	142	CD
17	72	262	CD
18	76	234	CD
20	10	89	CD
24	86	615	CD

^a Percent decrease in urine volume vs. control at a dose of 10 mg/kg

5 ^b Osmolality changes expressed as percent of control at a dose of 10 mg/kg

^c Rat model used: Sprague-Dawley (CD)

10 The following examples are presented to illustrate rather than limit the scope of the invention.

EXAMPLE 1**[2-Chloro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido [2,3-b] [1,5] benzodiazepin-6-yl)-methanone**

5

Step A. 6,11-Dihydro-5H-pyrido[2,3-b][1,5]benzodiazepin-5-one 1:1 salt with hydrochloric acid

10 A mixture of 1,2-phenylene diamine (52 g, 480 mmol) and chloronicotinic acid (76 g, 482 mmol) in cyclohexanol (480 mL) was refluxed under nitrogen for 2.5 hours. A precipitate appeared soon after the heating was initiated. The warm reaction mixture was carefully poured onto ice-cold dichloromethane (1000 mL) under vigorous stirring. The semisolid mass was collected, washed thoroughly with dichloromethane and dried in vacuo to yield 98.9 g (83%) of the title compound
15 which was used in the next step without further purification.

Step B. 6,11-Dihydro-5H-pyrido[2,3-b][1,5]benzodiazepine

20 Diborane dimethylsulfide complex (35 mL) was added via syringe to a suspension of 6,11-dihydro-5H-pyrido[2,3-b][1,5]benzodiazepin-5-one 1:1 salt with hydrochloric acid of Step A (25 g, 0.1 mole) in dioxane (230 mL) under nitrogen. The mixture was sonicated overnight at room temperature and then evaporated to dryness in vacuo. The green residue was treated with cold 2N hydrochloric acid and diethyl ether. The cold aqueous layer was basified with 50% aqueous sodium
25 hydroxide (to pH 9) and the basic layer extracted with ethyl acetate. The organic extracts were dried over anhydrous potassium carbonate, and evaporated to dryness to yield a burgundy solid (24.35 g, 61.4%). This crude material was purified by trituration with diethyl ether. The solid was collected, washed and dried in vacuo. The mother liquors from different runs were combined and the mixture (18.5 g) flash
30 chromatographed (on silica Merck-60, eluant 20% ethyl acetate in hexane) to provide additional material homogeneous by TLC (yellow solid, 11 g).

Step C. 2-Chloro-4-fluorobenzoyl chloride

A suspension of the 2-chloro-4-fluorobenzoic acid (13.61g, 78 mmol) in dichloromethane (85 mL) containing a few drops of dimethylformamide was treated dropwise under nitrogen with a 2M solution of oxalyl chloride in dichloromethane (1.2 equivalents). After gas evolution subsided, the reaction mixture was refluxed for an additional 25 minutes and then evaporated to dryness in vacuo. The crude acid chloride was used as such in the next step.

10 Step D. (2-Chloro-4-fluorophenyl)-(5,11-dihydro-pyrido [2,3-b] [1,5] benzodiazepin-6-yl)-methanone

To a solution of 6,11-dihydro-5H-pyrido[2,3-b][1,5]benzodiazepine of Step B (12.8 g, 65 mmol) in dimethylformamide (120 mL) under nitrogen was added potassium carbonate (19.76 g, 143 mmol). The mixture was cooled and treated dropwise with a solution of crude 2-chloro-4-fluorobenzoyl chloride of Step C (78 mmol) in dimethylformamide (50 mL). After stirring at room temperature for 75 minutes the mixture was diluted with water and extracted with dichloromethane. The organic extracts were dried over magnesium sulfate and evaporated to dryness. The crude material was purified by flash chromatography (on silica Merck-60, hexane-ethyl acetate gradient from 95:5 to 80:20) to provide the pure title compound (14.25 g, 62%) along with some less pure material (2.7 g). The pure material is an off-white crystalline solid, which is used as such in the next step.

NMR (DMSO- d_6 , 400 MHz): δ 4.13 and 5.42 (dd, 2 H, CONCH₂), 6.52 (m, 1H), 6.71-6.79 (m, 2H), 6.98-7.16 (2 m, 2H), 7.23-7.33 (m, 3H), 7.58 (m, 1H), 8.10 (m, 1H), 9.53 (s, 1H, NH)

MS (EI, m/z): 353/355 [M]⁺, 196

30 Step E. [2-Chloro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido [2,3-b] [1,5] benzodiazepin-6-yl)-methanone

Sodium hydride (60% suspension in oil, 1.8 g, 45.19 mmol) was washed with hexane, dried under nitrogen and resuspended in dry dimethylformamide (130 mL). Neat 3-methyl pyrazole (3.71 g, 45.19 mmol) was added dropwise at 0°C. After the

gas evolution subsided the cooling bath was removed and stirring was continued at room temperature. The (2-chloro-4-fluorophenyl)-(5,11-dihydro-pyrido[2,3-b][1,5] benzodiazepin-6-yl)- methanone of Step D (8.11 g, 22.59 mmol) was added in one portion and the mixture was placed in an oil bath (preheated at 130°C) for 2 hrs.

5 After cooling, the mixture was partitioned between water and ethyl acetate. The organic extracts were dried over sodium sulfate, and evaporated to dryness in vacuo. The residue was dissolved in dichloromethane and absorbed onto a silica Merck-60 flash column. Elution with a hexane-ethyl acetate gradient (from 95:5 to 3:2) provided the desired product along with some mixed fractions containing the title

10 compound and its more polar 5-methylpyrazole regioisomer of Example 2. The title compound crystallized by sonication from hexane-ethanol as a white solid (6.4 g, 68%), m.p. 207°C.

NMR (DMSO- d_6 , 400 MHz): δ 2.21 (s, 3H), 4.14 and 5.45 (dd, 2H, CONCH₂), 6.32 (m, 1H, pyrazole CH), 6.51 (m, 1H), 6.74-6.79 (2m, 2H), 6.98 (m, 1H), 7.25 (m, 2H),

15 7.58-7.70 (mm, 3H), 8.11 (m, 1H), 8.38 (m, 1H, pyrazole CH), 9.55 (s, 1H, NH)

MS (EI, m/z): 415/417 [M]⁺; (+FAB, m/z): 416/418 [M+H]⁺

Anal. Calc'd for C₂₃H₁₈ClN₅O: C 66.43; H 4.36; N 16.84. Found: C 66.11; H 4.42; N 16.64

20

EXAMPLE 2

[2-Chloro-4-(5-methyl-pyrazol-1-yl)-phenyl] (5,11-dihydro-pyrido [2,3-b] [1,5]benzodiazepin-6-yl)-methanone

25

The fractions (0.543 g) containing a mixture of 3-methyl and 5-methylpyrazole regioisomers obtained as described in Example 1, Step E were subjected to flash chromatography (silica gel Merck-60, eluant: toluene-ethyl acetate 90:10 followed by toluene-ethyl acetate-acetonitrile 90:10:5) to provide 0.327 g of the already described 3-methyl isomer of Example 1 and 0.105 g of the title compound as

30

an amorphous solid upon sonication from ether-hexane.

NMR (DMSO- d_6 , 400 MHz): δ 2.27 (s, 3H), 4.16 and 5.45 (dd, 2H, CONCH₂), 6.25 (m, 1H), 6.54 (m, 1H, pyrazole CH), 6.79 (m, 2H), 7.01 (m, 1H), 7.26 (m,

35

1H), 7.40-7.54 (mm, 3H), 7.61 (m, 1H), 8.11 (m, 1H, pyrazole CH), 9.56 (s, 1H, NH)

MS [EI, m/z]: 415/417 [M]⁺, 219/221, 196

EXAMPLE 3

5 [2-Bromo-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido [2,3-b] [1,5]benzodiazepin-6-yl)-methanone

Step A. 2-Bromo-4-fluorobenzoyl chloride

10 A suspension of 2-bromo-4-fluorobenzoic acid (6.87 g, 31.37 mmol) in dichloromethane (70 mL) containing a few drops of dimethylformamide was treated dropwise under nitrogen with a 2M solution of oxalyl chloride in dichloromethane (1.16 equivalents). After gas evolution subsided, the reaction mixture was refluxed for an additional 25 minutes and then the solution was evaporated to dryness in
15 vacuo. The crude acid chloride was used as such in the next step.

Step B. [2-Bromo-4-fluorophenyl]-(5,11-dihydro-pyrido [2,3-b] [1,5] benzodiazepin-6-yl)-methanone

20 To a solution of 6,11-dihydro-5H-pyrido[2,3-b][1,5]benzodiazepine of Example 1, Step B (5.15 g, 26.1 mmol) in dimethylformamide (70 mL) under nitrogen was added potassium carbonate (7.95 g, 57.51 mmol). The mixture was cooled and treated dropwise with a solution of crude 2-bromo-4-fluorobenzoyl chloride of Step A (31.37 mmol) in dimethylformamide (30 mL). After stirring at
25 room temperature for 75 minutes the mixture was diluted with water and extracted with dichloromethane. The organic extracts were dried over magnesium sulfate and evaporated to dryness to give a brown solid foam. The crude material was dissolved in dichloromethane and absorbed onto a silica Merck-60 flash column. Elution with a hexane-ethyl acetate gradient (from 95:5 to 75:25) provided the pure title compound
30 (6.18 g, 59.5%) along with some impure material (1.2 g). The pure material was triturated with hexane to provide an off-white solid foam, which was used as such in the next step.

-61-

NMR (DMSO- d_6 , 400 MHz): δ 4.13 and 5.42 (dd, 2 H, CONCH₂), 6.53 (m, 1H), 6.74-6.79 (m, 2H), 6.98-7.16 (2 m, 3H), 7.25 (m, 1H), 7.40-7.50 (broad s, 1H), 7.59 (m, 1H), 8.1 (m, 1H), 9.54 (s, 1H, NH)
MS (EI, m/z): 397/399 [M]⁺, 196

5

Step C. [2-Bromo-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido [2,3-b] [1,5]benzodiazepin-6-yl)-methanone

Sodium hydride (60% suspension in oil, 1.2 g, 30.15 mmol) was washed with
10 hexane, dried under nitrogen and resuspended in dry dimethylformamide (110 mL). Neat 3-methylpyrazole (2.47 g, 30.15 mmol) was added dropwise at 0°C. After the gas evolution subsided the cooling bath was removed and stirring was continued at room temperature. The [2-bromo-4-fluorophenyl]-(5,11-dihydro-pyrido [2,3-b] [1,5] benzodiazepin-6-yl)-methanone of Step B (6 g, 18.07 mmol) was added in one
15 portion to the clear solution. The mixture was placed in an oil bath (preheated at 130°C) for 40 minutes, cooled and partitioned between water and ethyl acetate. The organic extracts were dried over magnesium sulfate and evaporated to dryness. The crude material was dissolved in dichloromethane and absorbed onto a silica Merck-60
20 flash column. Elution with a hexane-ethyl acetate gradient (from 95:5 to 75:25) provided the less polar title compound (3.87 g) along with a mixture of 3- and 5-methylpyrazole regioisomers (0.860 g). The title compound (3.5 g, 51%) crystallized by sonication from hexane-ethanol, m.p. 208-209°C (dec):

NMR (DMSO- d_6 , 400 MHz): δ 2.21 (s, 3H), 4.15 and 5.44 (dd, 2H, CONCH₂), 6.31 (m, 1H, pyrazole CH), 6.52 (m, 1H), 6.77-6.80 (2m, 2H), 6.99 (m, 1H), 7.25 (m, 1H),
25 7.59-7.63 (2 m, 2H), 7.88 (m, 1H), 8.11 (m, 1H), 8.37 (s, 1H, pyrazole CH), 9.55 (s, 1H, NH)

MS (+EI, m/z): 459/461 [M]⁺, 265/263

Anal. Calc'd for C₂₃H₁₈BrN₅O: C 60.01, H 3.94, N 15.21. Found: C 59.92, H 4.05, N 15.01

30

EXAMPLE 4**(5,11-Dihydro-pyrido[2,3-b] [1,5]benzodiazepin-6-yl)-[4-(3-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-methanone****5 Step A. 2-Trifluoromethyl-4-fluorobenzoyl chloride**

A suspension of 2-trifluoromethyl-4-fluorobenzoic acid (16.85 g, 81 mmol) in dichloromethane (150 mL) containing a few drops of dimethylformamide was treated dropwise under nitrogen with oxalyl chloride (8.5 mL, 97.4 mmol). After the gas evolution subsided, the reaction mixture was refluxed for an additional 10 minutes, and then evaporated to dryness in vacuo. The crude acid chloride was used as such in the next step.

15 Step B. (5,11-Dihydro-pyrido[2,3-b] [1,5]benzodiazepin-6-yl)-(4-fluoro-2-trifluoromethyl-phenyl)-methanone

To a solution of 6,11-dihydro-5H-pyrido[2,3-b][1,5]benzodiazepine of Example 1, Step B (10.6 g, 53.8 mmol) in dimethylformamide (125 mL) under nitrogen was added potassium carbonate (22.4 g, 162 mmol). The mixture was cooled and treated dropwise with a solution of crude 2-trifluoromethyl-4-fluorobenzoyl chloride of Step A (81 mmol) in dimethylformamide (25 mL). After stirring at room temperature for 2 hours, the mixture was diluted with water and extracted with dichloromethane. The organic extracts were dried over magnesium sulfate and evaporated to dryness. The crude material was dissolved in dichloromethane and purified by flash chromatography (on silica Merck-60, hexane-ethyl acetate 80:20) to provide the pure title compound (6.9 g, 33.1%) which was crystallized by sonication from ethanol-hexane, m.p. 183-185°C.

NMR (DMSO-d₆, 400 MHz) δ 4.16 and 5.43 (dd, 2 H, CONCH₂), 6.56 (m, 1H), 6.64 (m, 1H), 6.79 (m, 1H), 7.02 (m, 1H), 7.26-7.40 (m, 3H), 7.58-7.65 (m, 2H), 8.12 (m, 1H), 9.59 (s, 1H, NH)

MS (EI, m/z): 387 [M]⁺

Anal. Calc'd for C₂₀H₁₃F₄N₃O: C 62.02, H 3.38, N 10.85. Found: C 62.06, H 3.22, N 10.67

Step C. (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[4-(3-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-methanone

Sodium hydride (60% suspension in oil, 0.83 g, 20.8 mmol) was washed once
5 with hexane, dried under nitrogen and resuspended in dry dimethylformamide (60 mL). 3-methyl pyrazole (0.90 mL, 11.2 mmol) was added in one portion. After the gas evolution subsided the stirring was continued at room temperature. The (5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-(4-fluoro-2-trifluoromethyl phenyl)-methanone of Step B (3.6 g, 9.3 mmol) was added in one portion and the mixture
10 was placed in an oil bath (preheated at 130°C) for 30 minutes. After cooling, the mixture was partitioned between water and ethyl acetate. The organic extracts were dried over sodium sulfate, and evaporated to dryness. The residue was dissolved in dichloromethane and absorbed onto a silica Merck-60 flash column. Elution with 25% ethyl acetate in hexane provided 3.3 g (79%) of the desired product as a foam
15 which crystallized by sonication from ethanol-hexane, m.p. 212-214°C. Further elution with 30% ethyl acetate in hexane yielded the more polar 5-methylpyrazole regioisomer of Example 6.

NMR (DMSO-d₆, 400 MHz): δ 2.23 (s, 3H, CH₃), 4.17 and 5.45 (dd, 2H, CONCH₂),
6.35 (m, 1H, pyrazole CH), 6.54 (m, 1H), 6.68 (m, 1H), 6.80 (m, 1H), 7.00 (m, 1H),
20 7.29 (m, 1H), 7.60 (m, 1H), 7.85 (m, 1H), 8.04 (m, 1H), 8.13 (m, 1H), 8.46 (m, 1H, pyrazole CH), 9.61 (s, 1H, NH)

MS (EI, m/z): 449 [M]⁺

Anal. Calc'd for C₂₄H₁₈F₃N₅O: C 64.14, H 4.04, N 15.58. Found: C 64.01, H 4.01, N 15.45

25

30

EXAMPLE 5**(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[4-(3-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-methanone**

5

Step A. 4-Fluoro-2-trifluoromethylbenzoic acid methyl ester

A suspension of 4-fluoro-2-trifluoromethylbenzoic acid (25.6 g, 123.0 mmol) in dichloromethane (250 mL) containing a few drops of dimethylformamide was treated dropwise under nitrogen with oxalyl chloride (11.3 mL, 129.5 mmol). After the gas evolution subsided, the reaction mixture was refluxed for an additional 15 minutes. The mixture was cooled and methanol (50 mL) was added. After stirring for 2 hrs, the reaction was concentrated, and the residue was partitioned between dichloromethane and water. The organic phase was washed with saturated aqueous sodium bicarbonate, dried over sodium sulfate, and evaporated to dryness to give 18.0 g (65.9%) of the title compound as a golden oil.

15

NMR (DMSO- d_6 , 400 MHz): δ 3.85 (s, 3H), 7.67 (m, 1H), 7.80 (m, 1H), 7.95 (m, 1H)MS (EI, m/z): 222 [M]⁺

The aqueous layer was acidified with 2 N hydrochloric acid and the white solid was collected by filtration to give 7.5 g (29.3%) of the starting 4-fluoro-2-trifluoromethylbenzoic acid.

20

Step B. 4-(3-Methyl-pyrazol-1-yl)-2-trifluoromethyl benzoic acid methyl ester

25

Sodium hydride (60% suspension in oil, 3.85 g, 96.3 mmol) was washed with hexane, dried under nitrogen and resuspended in dry dimethylformamide (150 mL). A solution of 3-methylpyrazole (7.75 mL, 96.3 mmol) in dimethylformamide (50 mL) was added dropwise at ambient temperature. Stirring was continued until the gas evolution subsided, and then a solution of methyl 4-fluoro-2-trifluoromethylbenzoic acid methylester of Step A (17.8 g, 80.1 mmol) in dimethylformamide (50 mL) was added dropwise to the clear solution. After stirring for 30 min at room temperature, the reaction was quenched with saturated aqueous ammonium chloride and extracted with

30

35

ethyl acetate. The organic extracts were dried over sodium sulfate and evaporated to dryness. The residue was dissolved in 1:1 mixture of dichloromethane and hexane and absorbed onto a silica Merck-60 flash column. Elution with a dichloromethane-hexane gradient (from 1:1 to 4:1) provided the title compound (13.6 g, 59.7%) as a white solid, m.p. 59-61°C.

NMR (DMSO-d₆, 400 MHz): δ 2.28 (s, 3H), 3.86 (s, 3H, CO₂CH₃), 6.43 (m, 1H, pyrazole CH), 7.97 (m, 1H), 8.18 (m, 1H), 8.23 (m, 1H), 8.62 (m, 1H, pyrazole CH)
MS (EI, m/z): 284 [M]⁺

Anal. Calc'd for C₁₃H₁₁F₃N₂O₂: C 54.93, H 3.90, N 9.86. Found: C 54.80, H 3.73
N 9.81

Step C. 4-(3-Methyl-pyrazol-1-yl)-2-trifluoromethyl benzoic acid

To a solution of 4-(3-methylpyrazol-1-yl)-2-trifluoromethylbenzoic acid methylester of Step B (1.19 g, 4.2 mmol) in methanol (10 mL) was added 2.5 N sodium hydroxide (3.3 mL, 8.3 mmol). The mixture was heated at reflux for 90 minutes, cooled and concentrated. The residue was partitioned between ethyl acetate and 1N hydrochloric acid. The organic extracts were dried over sodium sulfate and evaporated to dryness to give the title compound (1.14 g, quantitative yield) as a white solid, m.p. 192-194°C.

NMR (DMSO-d₆, 400 MHz): δ 2.28 (s, 3H), 6.42 (m, 1H, pyrazole CH), 7.95 (m, 1H), 8.14 (m, 1H), 8.20 (m, 1H), 8.61 (m, 1H, pyrazole CH), 13.4-13.7 (broad s, 1H, COOH)

MS (+FAB, m/z): 271 [M+H]⁺

Anal. Calc'd for C₁₂H₉F₃N₂O₂: C 53.34, H 3.36, N 10.37. Found: C 53.35, H 3.29, N 10.21

Step D. (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[4-(3-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-methanone

To a solution of 4-(3-methyl-pyrazol-1-yl)-2-trifluoromethyl benzoic acid (1.1 g, 4.1 mmole) of Step C, and triethylamine (0.57 mL, 4.1 mmol) in dichloromethane (20 mL) was added 1,3,5-trichlorobenzoyl chloride (0.63 mL, 4.0 mmol). After stirring for 5.5 hours, 6,11-dihydro-5H-pyrido[2,3-b][1,5]benzodiazepine of Example 1, Step B (0.67 g, 3.4 mmol) and 4-dimethylamino

pyridine (0.42 g, 3.4 mmol) were added. After stirring for an additional 18 hours, the mixture was poured into saturated aqueous sodium bicarbonate. The organic phase was washed with brine, dried over sodium sulfate, and evaporated to dryness. The residue was dissolved in dichloromethane and absorbed onto a silica Merck-60 flash column. Elution with hexane-ethyl acetate (gradient from 8:2 to 7:3) provided the title product (0.89 g, 58.2%) as a foam which crystallized by sonication from ethanol-hexane, m.p. 212-214°C. This material was identical to the compound of Example 4.

EXAMPLE 6

(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[4-(5-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-methanone solvate with 0.09 dichloromethane and 0.13 ethyl acetate

The title compound (0.350 g, 8%) was obtained as described in Example 4 above, as a foam which crystallized by sonication from ethanol-hexane, m.p. 238-240°C.

NMR (DMSO- d_6 , 400 MHz): δ 2.29 (s, 3H), 4.19 and 5.46 (dd, 2H, CONCH₂), 6.28 (m, 1H, pyrazole CH), 6.57 (m, 1H), 6.71 (m, 1H), 6.80 (m, 1H), 7.02 (m, 1H), 7.29 (m, 1H), 7.58-7.67 (m, 4H), 7.81 (m, 1H), 8.13 (m, 1H), 9.63 (s, 1H, NH)

MS (+FAB, m/z): 450 [M+H]⁺

Anal. Calc'd for C₂₄H₁₈F₃N₅O + 0.09 CH₂Cl₂ + 0.13 C₄H₈O₂: C 63.09, H 4.13, N 14.95. Found: C 63.39, H 4.23, N 14.89

EXAMPLE 7

(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[2-trifluoromethyl-4-(3-trifluoromethyl-pyrazol-1-yl)-phenyl]-methanone

Sodium hydride (60% suspension in oil, 0.17 g, 4.25 mmol) was washed with hexane, dried under nitrogen and resuspended in dry dimethylformamide (10 mL). 3-

-67-

trifluoromethyl pyrazole (0.34 g, 2.5 mmol) was added in one portion. After the gas evolution subsided stirring was continued at room temperature. The (5,11-dihydro-pyrido [2,3-b] [1,5] benzodiazepin-6-yl)- (4-fluoro-2-trifluoromethyl-phenyl)-methanone of Example 4, Step B (0.75 g, 1.94 mmol) was added in one portion and the mixture was placed in an oil bath (preheated at 130°C) overnight. After cooling, the mixture was partitioned between water and ethyl acetate. The organic extracts were dried over sodium sulfate, and evaporated to dryness in vacuo. The residue was crystallized from ethanol to yield the title compound (0.57 g, 57.3%) as an off-white solid, m.p. 127-129°C.

NMR (DMSO-d₆, 400 MHz): δ 4.19 and 5.46 (dd, 2H, CONCH₂), 6.54 (m, 1H), 6.70 (m, 1H), 6.80 (m, 1H), 7.02 (m, 1H), 7.07 (m, 1H, pyrazole CH), 7.29 (m, 1H), 7.61 (m, 1H), 8.00 (m, 1H), 8.05-8.16 (m, 2H), 8.84 (m, 1H, pyrazole CH), 9.63 (s, 1H, NH)

MS (EI, m/z): 503 [M]⁺

Anal. Calc'd for C₂₄H₁₅F₆N₅O: C 57.26, H 3.00, N 13.91. Found: C 57.07, H 2.97, N 13.58

EXAMPLE 8

(5-Methyl-5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[4-(3-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]methanone

Sodium hydride (60% suspension in oil, 0.12 g, 3.0 mmol) was washed with hexane, dried under nitrogen and resuspended in dry dimethylformamide (20 mL). Neat 3-methylpyrazole (0.13 mL, 1.61 mmol) was added dropwise at ambient temperature. Stirring was continued until the gas evolution subsided, and then the (5,11-dihydro-pyrido [2,3-b][1,5]benzodiazepin-6-yl)-(4-fluoro-2-trifluoromethyl-phenyl)methanone of Example 4, Step B (0.52 g, 1.34 mmol) was added in one portion to the clear solution. The mixture was placed in an oil bath (preheated at 130°C) for 30 minutes. After cooling, sodium hydride (60% suspension in oil, 0.080 g, 2.0 mmol, washed with hexane) and methyl iodide (0.25 mL, 4.0 mmol) were added. The reaction mixture was stirred for an additional 15 minutes, and then partitioned between water and ethyl acetate. The organic extracts were dried over sodium sulfate and evaporated to dryness. The residue was dissolved in dichloromethane and absorbed

onto a silica Merck-60 flash column. Elution with 20% ethyl acetate in hexane provided the title product (0.28 g, 45.1%) as a foam which crystallized by sonication from ethanol-hexane as a white solid, m.p. 188-190°C.

NMR (DMSO-d₆, 400 MHz): δ 2.23 (s, 3H, CCH₃), 3.55 (s, 3H, NCH₃), 4.37-4.43 (broad s, 1H, CONCH₂), 5.71-5.76 (broad s, 1H, CONCH₂), 6.35 (m, 1H, pyrazole CH), 6.89-6.93 (m, 3H), 7.19-7.24 (m, 2H), 7.31 (m, 1H), 7.61 (m, 1H), 7.90 (m, 1H), 8.06 (m, 1H), 8.24 (m, 1H), 8.48 (m, 1H, pyrazole CH)

MS (EI, m/z): 463 [M]⁺

Anal. Calc'd for C₂₅H₂₀F₃N₅O: C 64.79, H 4.35, N 15.11. Found: C 64.55, H 4.29, N 15.04

EXAMPLE 9

(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[2-fluoro-4-(3-methyl-pyrazol-1-yl)-phenyl]-methanone 0.19 hydrate

Step A. 2,4-Difluoro benzoylchloride

A suspension of 2,4-difluorobenzoic acid (3.6 g, 22.8 mmol) in dichloromethane (40 mL) containing a few drops of dimethylformamide was treated dropwise under nitrogen with oxalyl chloride (2.4 mL, 27.5 mmol). After gas evolution subsided, the reaction mixture was refluxed for an additional 15 minutes, and then the solution was evaporated to dryness in vacuo. The crude acid chloride was used as such in the next step.

Step B. (2,4-Difluoro-phenyl)-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone

To a solution of 5,11-dihydro-5H-pyrido[2,3-b][1,5]benzodiazepine of Example 1, Step B (3.0 g, 15.2 mmol) in dimethylformamide (35 mL) under nitrogen was added potassium carbonate (6.3 g, 45.6 mmol) followed by a solution of the crude 2,4-difluorobenzoylchloride of Step A (22.8 mmol) in dimethylformamide (15 mL). After stirring at room temperature for 20 minutes, the reaction mixture was washed with water and stirred to give a solid which was collected by filtration. The solid was

dissolved in chloroform and washed with 1 N sodium hydroxide and brine. The organic phase was dried over sodium sulfate and evaporated to dryness. The crude material was dissolved in dichloromethane and absorbed onto a silica Merck-60 flash column. Elution with 20% ethyl acetate in hexane provided the title compound (2.6 g, 51%) as a white foam which crystallized by sonication from hexane-ethanol, m.p. 161-163°C.

NMR (DMSO- d_6 , 400 MHz): δ 4.12-5.46 (dd, 2H, CONCH₂), 6.52 (m, 1H), 6.67 (m, 1H), 6.76 (m, 1H), 6.98-7.07 (m, 3H), 7.26 (m, 1H), 7.35 (m, 1H), 7.57 (m, 1H), 8.10 (m, 1H), 9.56 (s, 1H, NH)

MS (EI, m/z): 337 [M]⁺

Anal. Calc'd for C₁₉H₁₃F₂N₃O: C 67.65; H 3.88; N 12.46. Found: C 67.30; H 3.98; N 12.10

Step C. (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[2-fluoro-4-(3-methyl-pyrazol-1-yl)-phenyl]-methanone 0.19 hydrate

Sodium hydride (60% suspension in oil, 0.48 g, 12.0 mmol) was washed with hexane, dried under nitrogen and resuspended in dry dimethylformamide (60 mL). Neat 3-methylpyrazole (0.48 mL, 6.0 mmol) was added. Stirring was continued until the gas evolution subsided. The (2,4-difluoro-phenyl)-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone of Step B (2.0 g, 5.9 mmol) was added in one portion to the clear solution. The mixture was placed in an oil bath (preheated at 130°C) for 1 hour, cooled and partitioned between water and ethyl acetate. The organic extracts were dried over sodium sulfate and evaporated to dryness. The crude material was dissolved in dichloromethane and absorbed onto a silica Merck-60 flash column. Elution with hexane-ethyl acetate (gradient from 9:1 to 1:1) provided the title compound along with the more polar 4-fluoro regioisomer of Example 10. The title compound (0.30 g, 12.7%) was obtained as a foam which crystallized by sonication from hexane-ethanol, m.p. 122-125°C.

NMR (DMSO- d_6 , 400 MHz): δ 2.21 (s, 3H, CH₃), 4.13 and 5.48 (dd, 2H, CONCH₂), 6.32 (m, 1H, pyrazole CH), 6.51 (m, 1H), 6.70 (m, 1H), 6.77 (m, 1H), 7.01 (m, 1H), 7.27 (m, 1H), 7.35 (m, 1H), 7.41 (m, 1H), 7.53-7.59 (m, 2H), 8.10 (m, 1H), 8.35 (m, 1H, pyrazole CH), 9.57 (s, 1H, NH)

-70-

MS (EI, m/z): 399 [M]⁺

Anal. Calc'd for C₂₃H₁₈FN₅O + 0.19 H₂O: C 68.57, H 4.60, N 17.38. Found:

C 68.53, H 4.68, N 17.56

5

EXAMPLE 10

(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[4-fluoro-2-(3-methyl-pyrazol-1-yl)-phenyl]-methanone solvate with 0.20 ethanol

10

The title product was obtained along with its 2-fluoro regioisomer as described in Example 9. The material was further purified by preparative HPLC (Waters silica cartridge, 55:45 hexane-ethyl acetate as the eluant, flow rate 150 mL/min, detection at 254 nm) to give the pure title compound (0.25 g, 10.6%) as a foam which crystallized by sonication from hexane-ethanol, m.p. 180-181°C.

15

MS (EI, m/z): 399 [M]⁺

Anal. Calc'd for C₂₃H₁₈FN₅O + 0.20 C₂H₆O: C 68.78, H 4.74, N 17.14. Found:

C 68.67, H 4.76, N 16.97

EXAMPLE 11

20

[2-Chloro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5-methyl-5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone

Step A. (2-Chloro-4-fluoro-phenyl)-(5-methyl-5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone

25

Sodium hydride (60% suspension in oil, 0.76 g, 19 mmol) was washed once with hexane, dried under nitrogen and resuspended in dry dimethylformamide (10 mL). A solution of (2-chloro-4-fluorophenyl)-(5,11-dihydro-pyrido [2,3-b] [1,5] benzodiazepin-6-yl)-methanone of Example 1, Step D (6.12 g, 17.3 mmol) in dimethylformamide (50 mL) was added dropwise at 0°C. After the gas evolution subsided the cooling bath was removed and stirring was continued at room temperature. Methyl iodide (2.51 g, 17.7 mmol) was added in one portion to the yellow solution,

35 and stirring was continued for 1 hour. The mixture was partitioned between 20%

aqueous sodium chloride and dichloromethane. The organic extracts were dried over magnesium sulfate and evaporated to dryness. The residue was dissolved in dichloromethane and absorbed onto a silica Merck-60 flash column. Elution with
5 hexane-ethyl acetate gradient (from 95:5 to 85:15) provided the title product (4.55 g, 72%) as a white crystalline solid, m.p. 222-223°C.

NMR (DMSO- d_6 , 400 MHz): δ 3.46 (s, 3H, NCH₃), 4.37 and 5.67 (2 broad m, 2H, CONCH₂), 6.87-6.97 (m, 2H), 7.06-7.14 (m, 2H), 7.20-7.32 (m, 3H), 7.36 (m, 1H), 7.60 (m, 1H), 8.21 (m, 1H)

10 MS (EI, m/z): 367/369 [M]⁺; (+FAB, m/z): 368/370 [M+H]⁺

Anal. Calc'd for C₂₀H₁₅ClFN₃O: C 65.31, H 4.11, N 11.42. Found: C 65.04, H 4.14, N, 11.27

15 **Step B. [2-Chloro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5-methyl-5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone**

Sodium hydride (60% suspension in oil, 0.264 g, 6.6 mmol) was washed with hexane, dried under nitrogen and resuspended in dry dimethylformamide (8 mL). Neat 3-methylpyrazole (0.541 g, 6.6 mmol) was added dropwise at 0°C. After the gas
20 evolution subsided the cooling bath was removed and stirring was continued at room temperature. The (2-chloro-4-fluoro-phenyl)-(5-methyl-5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone of Step A (1.21 g, 3.3 mmol) was added in one portion to the clear solution. The mixture was placed in an oil bath (preheated at 130°C) for 30 minutes, cooled and partitioned between 20% aqueous sodium chloride
25 and dichloromethane. The organic extracts were dried over magnesium sulfate and evaporated to dryness. The residue was dissolved in dichloromethane and absorbed onto a silica Merck-60 flash column. Elution with hexane-ethyl acetate (gradient from 95:5 to 70:30) provided the less polar title compound (0.936 g, 66%) along with some mixture of 3- and 5-methylpyrazole regioisomers (0.220 g). The title
30 compound crystallized by sonication from hexane- ethanol as a white solid, m.p. 218-219°C (dec).

NMR (DMSO- d_6 , 400 MHz): δ 2.21 (s, 3H, pyrazole C-CH₃), 3.49 (s, 3H, NCH₃), 4.39 and 5.69 (broad dd, 2H, CONCH₂), 6.32 (m, 1H, pyrazole CH), 6.88-6.94 (mm, 2H), 7.10 (m, 1H), 7.18-7.24 (m, 1H), 7.26-7.30 (mm, 2H), 7.59-7.65 (m m, 2H),
35 7.77 (m, 1H), 8.22 (m, 1H), 8.38 (s, 1H, pyrazole CH)

MS (EI, m/z): 429/431 [M]⁺, 219, 195

Anal. Calc'd for C₂₄H₂₀ClN₃O: C 67.05, H 4.69, N 16.29. Found: C 67.26, H 4.69,

5 N 16.15

EXAMPLE 12

10 (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[2-methyl-5-(3-methyl-
pyrazol-1-yl)-phenyl]-methanone

Step A. 5-Fluoro-2-methyl benzoylchloride

15 A suspension of 5-fluoro-2-methyl benzoic acid (2.31 g, 15.0 mmol) in dichloromethane (30 mL) containing a few drops of dimethylformamide was treated dropwise under nitrogen with oxalyl chloride (1.6 mL, 18.3 mmol). After gas evolution subsided, the reaction mixture was refluxed for an additional 10 minutes, and then evaporated to dryness. The crude acid chloride was used as such in the next step.

20

Step B. (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-(5-fluoro-2-methyl-phenyl)-methanone

25 To a solution of 6,11-dihydro-5H-pyrido[2,3-b][1,5]benzodiazepine of Example 1, Step B (2.0 g, 10.1 mmol) in dimethylformamide (15 mL) under nitrogen was added potassium carbonate (4.1 g, 29.7 mmol). The mixture was treated dropwise with a solution of crude 5-fluoro-2-methyl benzoyl chloride of Step A (15.0 mmol) in dimethylformamide (10 mL). After stirring at room temperature for 15 minutes, the mixture was diluted with water and stirred to give a solid mass which
30 was collected by filtration. The solid was dissolved in chloroform and washed with 1 N sodium hydroxide and brine. The organic layer was dried over sodium sulfate and evaporated to dryness to give a purple oil. The crude material was dissolved in dichloromethane and absorbed onto a silica Merck-60 flash column. Elution with 20% ethyl acetate in hexane provided 1.88 g (55.8%) of the title product as a foam
35 which was crystallized by sonication from ethanol-hexane, m.p. 138-140°C.

-73-

NMR (DMSO- d_6 , 400 MHz): δ 1.95 (s, 3H, CH₃), 4.11 and 5.46 (dd, 2H, CONCH₂), 6.53 (m, 1H), 6.75-6.80 (m, 2H), 6.81-7.06 (m, 4H), 7.24 (m, 1H), 7.60 (m, 1H), 8.11 (m, 1H), 9.57 (s, 1H, NH)

MS (EI, m/z): 333 [M]⁺

- 5 Anal. Calc'd for C₂₀H₁₆FN₃O: C 72.06, H 4.84, N 12.60. Found: C 71.88, H 4.78, N 12.67

Step C. (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[2-methyl-5-(3-methyl-pyrazol-1-yl)-phenyl]-methanone

10

Sodium hydride (60% suspension in oil, 0.25 g, 6.25 mmol) was washed with hexane, dried under nitrogen and resuspended in dry dimethylformamide (10 mL). Neat 3-methylpyrazole (0.28 mL, 3.5 mmol) was added in one portion at ambient temperature. Stirring was continued until the gas evolution subsided. The 5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-(5-fluoro-2-methyl-phenyl)-methanone of Step B (0.75 g, 1.94 mmol) was added in one portion to the clear solution. The mixture was heated to reflux for 26 hours, cooled and partitioned between water and ethyl acetate. The organic extracts were dried over sodium sulfate and evaporated to dryness. The residue was dissolved in dichloromethane and absorbed onto a silica Merck-60 flash column. Elution with hexane-ethyl acetate (gradient from 8:2 to 7:3) provided the title product (0.55 g, 51.5%) as a pale yellow foam which was crystallized by sonication from hexane-ethanol, m.p. 209-210°C.

15

20

25

NMR (DMSO- d_6 , 400 MHz): δ 1.94 (s, 3H, CH₃), 2.23 (s, 3H, pyrazole CH₃), 4.13 and 5.49 (dd, 2H, CONCH₂), 6.28 (m, 1H, pyrazole CH), 6.50 (m, 1H), 6.78 (m, 2H), 6.97 (m, 1H), 7.07 (m, 1H), 7.24 (m, 1H), 7.51 (m, 1H), 7.62 (m, 1H), 8.11 (m, 1H, pyrazole CH), 8.19 (m, 1H), 9.60 (s, 1H, NH)

MS (EI, m/z): 395 [M]⁺

Anal. Calc'd for C₂₄H₂₁N₅O: C 72.89, H 5.35, N 17.71. Found: C 72.57, H 5.49, N 17.46

30

EXAMPLE 13**[4-(3-tert-Butyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone**

5 Sodium hydride (60% suspension in oil, 0.12 g, 3.0 mmol) was washed with hexane, dried under nitrogen and resuspended in dry dimethylformamide (10 mL). 3-tert-butylpyrazole (0.20 g, 1.6 mmol) was added in one portion at ambient temperature, and the stirring was continued until the gas evolution subsided. The
10 (5,11-dihydro-pyrido [2,3-b] [1,5] benzodiazepin-6-yl)- [4-fluoro-2-trifluoromethyl-phenyl]-methanone of Example 4, Step B (0.50 g, 1.3 mmol) was added in one portion to the clear solution. The mixture was placed in an oil bath (preheated at 130°C) for 30 minutes and then heated at reflux for 5 hours. After cooling, the mixture was partitioned between water and ethyl acetate. The organic extracts were
15 dried over sodium sulfate and evaporated to dryness. The crude residue was dissolved in dichloromethane and absorbed onto a silica Merck-60 flash column. Elution with 25% ethyl acetate in hexane provided the title product (0.23 g, 36.0%) as a foam which crystallized by trituration with hexane-ether, m.p. 136-140°C.
NMR (DMSO-d₆, 400 MHz): δ 1.26 (s, 9H, C(CH₃)₃), 4.17 and 5.45 (dd, 2H, CONCH₂), 6.47 (m, 1H, pyrazole CH), 6.54 (m, 1H), 6.68 (m, 1H), 6.80 (m, 1H),
20 7.00 (m, 1H), 7.28 (m, 1H), 7.60 (m, 1H), 7.87 (m, 1H), 8.04 (m, 1H), 8.13 (m, 1H), 8.47 (m, 1H, pyrazole CH), 9.62 (s, 1H, NH)
MS (EI, m/z): 491 [M]⁺
Anal. Calc'd for C₂₇H₂₄F₃N₅O: C 65.98, H 4.92, N 14.25. Found: C 65.75, H 4.92,
25 N 13.95

EXAMPLE 14

(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[4-(3-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-methanone 1:1 salt with methanesulfonic acid solvate with 0.15 hexane

30

To a solution of (5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[2-trifluoromethyl-4-(3-methyl-pyrazol-1-yl)-phenyl]-methanone of Example 4, Step C
35 (0.51 g, 1.13 mmol) in dichloromethane (10 mL) was added methanesulfonic acid

-75-

(0.074 mL, 1.14 mmol). After stirring overnight at room temperature, hexane was added and the precipitate was collected by filtration to give the title salt (0.54 g, 87.6%) as a white solid, m.p. 256-257°C.

MS (+FAB, m/z): 450 [M+H]⁺

- 5 Anal. Calc'd for C₂₄H₁₈F₃N₅O + CH₄O₃S + 0.15 C₆H₁₄: C 55.70, H 4.35, N 12.54.
Found: C 55.37, H 4.48, N 12.45

EXAMPLE 15

- 10 (5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[4-(3-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-methanone 1:1 salt with hydrochloric acid 0.10 dichloromethane solvate

- 15 To a solution of (5,11-dihydro-pyrido [2,3-b] [1,5] benzodiazepin-6-yl)-[2-trifluoromethyl-4-(3-methyl-pyrazol-1-yl)-phenyl]-methanone of Example 4, Step C (0.55 g, 1.22 mmol) in dichloromethane (10 mL) was added 1N hydrochloric acid in ether (1.22 mL). The mixture was stirred for 1 hour and the precipitate was collected by filtration to give the title salt (0.50 g, 84.1%) as an amorphous white solid.

MS (EI, m/z): 449 [M]⁺

- 20 Anal. Calc'd for C₂₄H₁₈F₃N₅O + 1.04 HCl + 0.10 CH₂Cl₂: C 58.38, H 3.91, N 14.12.
Found: C 58.23, H 4.07, N 14.03

EXAMPLE 16

- 25 [2-Chloro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(11H-5-oxa-4,10-diazadibenzo[a,d]cyclohepten-10-yl)-methanone

Step A. 6H-Pyrido[2,3-b][1,5]benzoxazepine

- 30 Under an atmosphere of nitrogen, lithium aluminum hydride (0.258 g, 6.877 mmol) was added portionwise under stirring to 6H-pyrido[2,3-b][1,5]benzoxazepin-5-one (0.636 g, 3 mmol) in 25 mL of tetrahydrofuran. After heating at reflux for 3 hours, excess lithium aluminum hydride was decomposed by the stepwise addition of 0.3 mL of water, 0.3 mL of 1N sodium hydroxide, 0.9 mL of water, and 3.8 g sodium sulfate. The solids were filtered and washed with ethyl acetate. The filtrate
35 was evaporated under reduced pressure to give crude product (0.45 g) which was

purified by flash chromatography (on silica Merck 60, dichloromethane-ethyl acetate, 19:1). The title compound was obtained as a yellow solid (0.10 g, 17 %).

NMR (DMSO- d_6 , 400 MHz): δ 4.32 (d, 2H, CH_2N), 6.09 (t, 1H, NH), 6.56 (t, 1H), 6.62 (d, 1H), 6.82 (t, 1H), 7.00 (d, 1H), 7.18 (m, 1H), 7.79 (d, 1H), 8.165 (m 1H).

5 MS (EI, m/z): 198 $[M]^+$

Step B. (2-Chloro-4-fluoro-phenyl)-(11H-5-oxa-4,10-diaza-dibenzo[a,d]cyclohepten-10-yl)-methanone

10 Under anhydrous conditions a mixture of 2-chloro-4-fluorobenzoic acid (0.37 g, 2.12 mmol) and oxalyl chloride (0.303 mL, 2.3 mmol) in 10 mL of dichloromethane containing a catalytic amount of dimethylformamide was stirred at room temperature until gas evolution ceased. The crude acid chloride solution was then added portionwise to a stirred solution of 6H-pyrido[2,3-b][1,5]benzoxazepine
15 of Step A (0.45 g, 2.12 mmol) and triethylamine (0.35 mL, 2.5 mmol) in 15 mL of dichloromethane. The reaction mixture was stirred overnight at ambient temperature, washed with water and dried over sodium sulfate. Evaporation of the solvent provided the crude title compound as a gummy solid which was used as such in the next step.

20 NMR (DMSO- d_6 , 400 MHz): δ 5.1 (broad, 2H, CH_2N), 6.9-7.5 (m, 8H), 7.92 (d, 1H), 8.22 (m, 1H)

MS (EI, m/z): 354 $[M]^+$, 319, 198, 157

Step C. [2-Chloro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(11H-5-oxa-4,10-diaza-dibenzo[a,d]cyclohepten-10-yl)-methanone

25 Under an atmosphere of nitrogen, 3-methylpyrazole (0.161 mL, 2.0 mmol) was added to a slurry of hexane washed potassium hydride (0.08g, 2.0 mmol) in 5 mL of
30 dry dimethylformamide. The mixture was stirred at ambient temperature until gas evolution ceased. A solution of 2-chloro-4-fluoro-phenyl-(11H-5-oxa-4,10-diaza-dibenzo[a,d]cyclohepten-10-yl)-methanone of Step B (0.79 g, 2.14 mmol) in 10 mL of
dimethylformamide was added. The mixture was heated at 130°C for 3.5 hours,
35 cooled and partitioned between ethyl acetate and brine. The organic phase was

-77-

washed with water, dried over sodium sulfate and concentrated to dryness in vacuo. The crude product was purified by flash chromatography (on silica Merck 60, dichloromethane-ethyl acetate gradient from 19:1 to 9:1) to give the title compound as a white solid (0.18 g, 21%), m.p. 220-222°C.

5

NMR (DMSO- d_6 , 400 MHz): δ 2.218 (s, 3H, CH₃), 5.106 (broad 2H, CH₂N), 6.325 (d, 1H), 6.927 (t, 1H), 7.03 (d, 1H), 7.21 (m, 2H), 7.27 (d, 1H), 7.44 (m, 1H), 7.67 (d, 1H), 7.70 (s, 1H), 7.88, (d, 1H), 8.24 (m, 1H), 8.397 (d, 1H).

MS (EI, m/z): 416 [M]⁺, 219

10

EXAMPLE 17

[2-Chloro-4-(3-trifluoromethyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-methanone

15

Sodium hydride (60% suspension in oil, 0.195 g) was washed with hexane, dried under nitrogen and resuspended in dry dimethylformamide (10 mL). 3-trifluoromethyl pyrazole (0.364 g) was added dropwise at 0°C. After the gas evolution subsided the solution was brought to room temperature. The (2-chloro-4-fluorophenyl)- (5,11-dihydro-pyrido [2,3-b] [1,5] benzodiazepin-6-y)-methanone of Example 1, Step D (0.787 g, 2.23 mmol) was added in one portion and the mixture was placed in an oil bath (preheated at 130°C) for 4.5 hours. The mixture was cooled and partitioned between saturated aqueous ammonium chloride and ethyl acetate. The organic extracts were dried over sodium sulfate and evaporated to dryness in vacuo. The residue was dissolved in dichloromethane and absorbed onto a silica Merck-60 flash column. Elution with hexane-ethyl acetate (gradient from 95:5 to 3:2) provided the desired product (0.727 g, 69%) which crystallized by sonication from hexane-ethanol as an off-white solid, m.p. 183-185°C.

20

25

30

NMR (DMSO- d_6 , 400 MHz): δ 4.16 and 5.45 (dd, 2H, CONCH₂), 6.52 (m, 1H), 6.78 (m, 2H), 7.01 (m, 2H), 7.04 (m, 1H, pyrazole CH), 7.26 (m, 1H), 7.61 (m,

1H), 7.74-7.84 (2m, 2H), 8.12 (m, 1H), 8.74 (m, 1H, pyrazole CH), 9.58 (s, 1H, NH)
MS (EI, m/z): 469/471 [M]⁺, 273/275, 196

Anal. Calc'd for $C_{23}H_{15}ClF_3N_5O$: C 58.80, H 3.22, N, 14.91. Found: C 58.67, H 3.14, N 14.83

EXAMPLE 18

5 **[2-Chloro-4-(1-methyl-1-H-pyrazol-3-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone**

Step A. 2-Chloro-4-(3-dimethylaminopropyn-1-yl)-benzoic acid methyl ester

- 10 Under an atmosphere of nitrogen, a mixture of 4-bromo-2-chlorobenzoic acid methyl ester (25.13 g, 101 mmol), 1-dimethylamino-2-propyne (16 mL, 150 mmol), bis(triphenylphosphine)palladium(II) chloride (1.0 g) and copper (I) iodide (0.15 g) in 100 mL of triethylamine was heated at 60°C for 2 hours. The cooled reaction mixture was filtered through Solka floc and the cake was washed with ethyl acetate.
- 15 The filtrate was partitioned between ethyl acetate and dilute aqueous sodium thiosulfate. The organic layer was washed with water, brine and dried over sodium sulfate. The dark solution was filtered through a plug of Merck-60 silica gel and the filtrate was concentrated in vacuo to give the title compound (23.8 g, 95%) as an orange oil, which was used as such in the next step.
- 20 NMR (DMSO- d_6 , 300 MHz): δ 2.25 (s, 6H, NCH_3), 3.475 (s, 2H, CCH_2N), 3.84 (s, 3H, OCH_3), 7.5 (dd, 1H), 7.62 (s, 1H), 7.8 (d, 1H).

Step B. 2-Chloro-4-(3-dimethylamino-2-propen-1-yl)-benzoic acid methyl ester

- 25 Under an atmosphere of nitrogen, purified meta-chloroperbenzoic acid (16.0 g, 93 mmol) was added portionwise to a stirred solution of 2-chloro-4-(3-dimethylaminopropyn-1-yl)-benzoic acid methyl ester of Step A (23.5 g, 93.4 mmol) in 200 mL of dichloromethane at -10°C. After the addition was complete, the solution was stirred at reduced temperature for 30 minutes and then filtered through a
- 30 column of

- basic alumina (400 g, Brockman activity I) packed with dichloromethane-methanol (9:1, v/v). The intermediate N-oxide was eluted with the above solvent system. The
- 35 dichloromethane was then carefully replaced with methanol by evaporation at or

below room temperature, taking care that the mixture is never allowed to evaporate to dryness. The methanolic solution was heated at 60°C overnight, and then was concentrated in vacuo. The residue was purified by flash chromatography (on Merck-60 silica gel, hexane-ethyl acetate 1:1) to give 12.1 g of a slightly impure product.

- 5 Trituration with diethyl ether provided the pure title compound (6.15 g, 48%) as an orange solid.

NMR (DMSO- d_6 , 300 MHz): δ 2.98 (s, 3H, NCH₃), 3.2 (s, 3H, NCH₃), 3.83 (s, 3H, OCH₃), 5.85 (d, 1H, vinyl H), 7.75-8.0 (m, 4H, vinyl H + ArH).

10 **Step C. 2-Chloro-4-(1H-pyrazol-3-yl)-benzoic acid methyl ester**

A solution of 2-chloro-4-(3-dimethylamino-2-propen-1-yl)-benzoic acid methyl ester of Step B (6.13 g, 22.9 mmol) and anhydrous hydrazine (1.44 mL, 45.8 mmol) in 15 mL of glacial acetic acid was heated at 90°C for 30 minutes. The

15 reaction mixture was concentrated in vacuo and the residue partitioned between ethyl acetate and water. The organic layer was washed with water and brine and dried over sodium sulfate. The solvent was evaporated and the residual solid triturated with diethyl ether-hexane to give the title compound (5.1g, 94%) as an orange solid.

- 20 NMR (DMSO- d_6 , 300 MHz): δ 3.85 (s, 3H, OCH₃), 6.9 (d, 1H), 7.9 (m, 3H), 8.0 (d, 1H), 13.15 (broad, 1H, NH).

Step D. 2-Chloro-4-(1-methyl-1H-pyrazol-3-yl)-benzoic acid methyl ester

- 25 Under an atmosphere of nitrogen, a solution of 2-chloro-4-(1H-pyrazol-3-yl)-benzoic acid methyl ester of Step C (5.0 g, 21.1 mmol) in 50 mL of dry dimethylformamide was added dropwise to a stirred mixture of hexane washed sodium hydride (0.51 g, 21.1 mmol) in 5 mL of dry dimethylformamide. The mixture was stirred at ambient temperature for 30 minutes, methyl iodide (2.7 mL, 42.2 mmol) was added to the resulting solution and the stirring was continued
- 30 overnight at room temperature. The reaction mixture was poured into water and extracted with ethyl

acetate. The organic layer was washed with water, brine and dried over sodium sulfate. Removal of solvent in vacuo afforded 4.8 g of an orange oil. Flash

chromatography of the crude material (on silica gel Merck-60, hexane-ethyl acetate, 4:1) provided 2.9 g (55%) of the desired 1-methylpyrazole regioisomer.

NMR (DMSO- d_6 , 300 MHz): δ 3.84 (s, 3H, CH₃), 3.9 (s, 3H CH₃), 6.875 (d, 1H), 7.8 (d, 1H), 7.85 (s, 2H), 7.95 (s, 1H).

5

Step E. 2-Chloro-4-(1-methyl-1H-pyrazol-3-yl)-benzoic acid

A solution of 2-chloro-4-(1-methyl-1H-pyrazol-3-yl)-benzoic acid methyl ester of Step D (2.9 g, 11.6 mmol) in 20 mL of methanol containing 5 mL of 2.5 N sodium hydroxide was stirred at ambient temperature overnight. An additional 2.0 mL of 2.5 N sodium hydroxide were added and the solution was gently heated for 30 minutes. The reaction mixture was concentrated in vacuo, diluted with water, and acidified with 2N hydrochloric acid. The precipitate was collected and thoroughly dried to give 2.55 g (93%) of the title compound.

15 NMR (DMSO- d_6 , 300 MHz): δ 3.9 (s, 3H, NCH₃), 6.85 (d, 1H), 7.82 (m, 3H), 7.95 (s, 1H), 13.3 (broad, 1H, COOH).

Step F. [2-Chloro-4-(1-methyl-1H-pyrazol-3-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone

20

Under anhydrous conditions a solution of 2-chloro-4-(1-methyl-1H-pyrazol-3-yl)-benzoic acid of Step E (2.1 g, 8.88 mmol) and triethylamine (1.3 mL, 9.2 mmol) in 75 mL of dichloromethane was treated in one portion with 2,4,6-trichlorobenzoyl chloride (1.48 mL, 9.2 mmol) and stirred at ambient temperature for 2 hours. To the solution was added the 6,11-dihydro-5H-pyrido[2,3-b][1,5]benzodiazepine of Example 1, Step B (1.74 g, 8.9 mmol) followed by 4-dimethylaminopyridine (1.1 g, 8.9 mmol) and stirring was continued for 18 hours. The reaction mixture was washed

25

sequentially with saturated sodium bicarbonate and brine. After drying over sodium sulfate, the solution was concentrated to small volume and absorbed onto silica Merck-60. Elution with ethyl acetate-hexane (gradient from 4:3 to 2:1) gave the pure title

30

compound as a syrup which crystallized from diethylether. The white solid (0.78 g, 23% yield based on recovered starting material) melted at 196-197°C.

NMR (DMSO-d₆, 400 Mhz): δ 3.831 (s, 3H, NCH₃), 4.13 (d, 1H), 5.43 (d, 1H), 6.497 (t, 1H), 6.71 (d, 1H), 6.76 (m, 2H), 6.97 (t, 1H), 7.24 (d, 1H), 7.6 (m, 3H), 7.705 (d, 1H), 8.10(dd, 1H), 9.544 (s, 1H, NH).

MS (EI, m/z): 415/417 [M]⁺, 219/221

Anal. Calc'd for C₂₃H₁₈ClN₅O: C 66.42, H 4.36, N 16.84. Found: C 66.20, H 4.49, N 16.59.

EXAMPLE 19

[2-Chloro-4-(1-methyl-1H-pyrazol-3-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone

Under anhydrous conditions a mixture of the 2-chloro-4-(1-methyl-1H-pyrazol-3-yl)-benzoic acid of Example 18, Step E (1.9 g, 8.05 mmol) and oxalyl chloride (0.79 mL, 9.0 mmol) in 20 mL of dichloromethane containing a catalytic amount of dimethylformamide was stirred at ambient temperature for 1 hour. The solvent was evaporated and the solid acid chloride was dissolved in 5 mL of dimethylformamide and added directly to a mixture of 6,11-dihydro-5H-pyrido[2,3-b][1,5]benzodiazepine of Example 1, Step B (1.59 g, 8.05 mmol) and potassium carbonate (1.25 g, 9.0 mmol). After stirring for 2 hours at ambient temperature the reaction mixture was partitioned between ethyl acetate and water. The organic phase was washed with water and brine, dried over sodium sulfate, and concentrated to small volume. Flash chromatography of the residue (on silica Merck 60, ethyl acetate-hexane, gradient from 4:3 to 2:1) gave the product as a syrup which crystallized from diethyl ether (1.8 g, 61% yield, based on recovered starting material) as a white solid, m.p. 196-197°C. Another recrystallization from ethanol-diethyl ether provided a higher melting polymorph, m.p. 202°C as determined by differential scanning calorimetry.

MS (+FAB, m/z): 416/418 (M+ H)⁺.

Anal. Calc'd for C₂₃H₁₈ClN₅O: C 66.42, H 4.36, N 16.84. Found: C 66.20, H 4.42, N 16.80.

EXAMPLE 20**[2-Chloro-4-(1-methyl-1H-pyrazol-3-yl)-phenyl]-(5-methyl-5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone**

5

Under anhydrous conditions, a solution of [2-chloro-4-(1-methyl-1H-pyrazol-3-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5] benzodiazepin-10-yl)-methanone of Example 19 (0.382 g, 0.92 mmol) in 10 mL of tetrahydrofuran was added dropwise to a stirred slurry of hexane washed sodium hydride (0.025g, 1.02 mmol) in 2 mL of tetrahydrofuran. After the gas evolution ceased, methyl iodide (1 mL) was added and stirring was continued for 2 hours. The reaction mixture was partitioned between ethyl acetate and water. The organic layer was washed with brine and dried over sodium sulfate. The solvent was removed and the crude product purified by flash chromatography (on silica Merck-60, dichloromethane-ethyl acetate 2:1) to give the title compound (0.18 g, 47%). Crystallization from diethylether provided a light yellow solid (0.16 g), m.p. 249-250°C.

15

NMR (DMSO-d₆, 400 MHz): δ 3.491 (s, 3H NCH₃), 3.835 (s, 3H, NCH₃), 4.18 (broad, 1H), 5.7 (broad, 1H), 6.497 (t, 1H), 6.72 (d, 1H), 6.88 (m, 2H), 7.08 (d, 1H), 7.19 (m, 2H), 7.25 (d, 1H), 7.6 (m, 2H), 7.69 (d, 1H), 7.71 (d, 1H), 8.215 (dd, 1H).

20 MS (EI, m/z): 429/431 [M]⁺, 219/221.

EXAMPLE 21**[2-Chloro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido [2,3-b][1,5] benzodiazepin-6-yl)-methanone**

25

Step A. 2-Chloro-4-(3-methyl-1H-pyrazol-1-yl)-benzoic acid methyl ester

Under anhydrous conditions a stirred suspension of hexane washed potassium hydride (0.424 g, 10.6 mmol) in 5 mL of dimethylformamide was treated in one portion with 3-methyl pyrazole (0.85 mL, 10.6 mmol). After the gas evolution ceased, 2-chloro-4-fluorobenzoic acid methyl ester (2.0 g, 10.6 mmol) was added to the clear

30

- solution. The mixture was heated at 130°C for 15 minutes, cooled, and partitioned between ethyl acetate and brine. The organic layer was washed with water and brine, and dried over sodium sulfate. Removal of solvent afforded 2.2 g of a yellow oil consisting of a mixture of 3-methyl and 5-methylpyrazole regioisomers. In addition, about 20% of the acid derived from hydrolysis of the ester was detected by analysis of the NMR spectrum of the crude product. The desired 3-methylpyrazole regioisomer was separated from the 5-methyl isomer of Example 22 by flash chromatography (on silica Merck-60, dichloromethane-hexane 2:1) and was isolated as a white solid (1.55 g, 56%).
- 10 NMR (DMSO- d_6 , 400 MHz): δ 2.264 (s, 3H, CCH₃), 3.845 (s, 3H, OCH₃), 6.40 (d, 1H), 7.865 (dd, 1H), 7.93 (d, 1H), 8.00 (s, 1H), 8.535 (d, 1H).
MS (EI, m/z): 250/252 [M]⁺, 219

Step B. 2-Chloro-4-(3-methyl-1H-pyrazol-1-yl)-benzoic acid

15

- A solution of 2-chloro-4-(3-methyl-1H-pyrazol-1-yl)-benzoic acid methyl ester of Step A (1.42 g, 5.6 mmol) in 20 mL of tetrahydrofuran containing 6 mL of 1 M aqueous lithium hydroxide was stirred overnight at ambient temperature. The reaction mixture was partitioned between ethyl acetate and 1 N hydrochloric acid. The organic layer was washed with water and brine, and dried over sodium sulfate. Evaporation of the solvent afforded the title compound (1.05 g, 78%), m.p. 192-193°C.
- 20 NMR (DMSO- d_6 , 400 MHz): δ 2.268 (s, 3H, CCH₃), 6.40 (d, 1H), 7.84 (dd, 1H), 7.92 (d, 1H), 8.00 (s, 1H), 8.53 (d, 1H), 13.32 (broad, 1H, COOH).
25 MS (EI, m/z): 236/238 [M]⁺, 219
Anal. Calc'd for C₁₁H₉ClN₂O₂: C, 55.83, H 3.83, N 11.84. Found: C 55.79, H 3.98, N 11.73

- Step C. [2-Chloro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-methanone**
- 30

In the manner of Example 5, Step D, employing 2-chloro-4-(3-methyl-1H-pyrazol-1-yl)-benzoic acid of Step B (0.971 g, 4.1 mmol), triethylamine (0.57 mL, 4.1 mmol), 1,3,5-trichlorobenzoylchloride (0.63 mL, 4.0 mmol), 6,11-dihydro-5H-

pyrido[2,3-b][1,5]benzodiazepine of Example 1, Step B (0.67 g, 3.4 mmol) and 4-dimethylamino pyridine (0.42 g, 3.4 mmol) in dichloromethane (20 mL), was obtained a compound identical to that of Example 1.

5

EXAMPLE 22**2-Chloro-4-(5-methyl-1H-pyrazol-1-yl)-benzoic acid methyl ester.**

The title compound was prepared as described in Example 21, Step A and separated from the 3-methylpyrazole isomer of Example 21 by flash chromatography (on silica Merck-60, eluant: dichloromethane). It was obtained as a white solid (0.20 g, 7.5 %).

10

NMR (DMSO-d₆, 400 Mhz): δ 2.425 (s, 3H, CCH₃), 3.875 (s, 3H, OCH₃), 6.33 (s, 1H), 7.65 (m, 2H), 7.79 (s, 1H), 7.95 (d, 1H).

MS (EI, m/z): 250/252 [M]⁺, 219.

15

EXAMPLE 23**(2-Chloro-4-fluorophenyl)-(5,11-dihydro-pyrido[2,3-b][1,5] benzodiazepin-6-yl-methanone**

20

To a solution of 6,11-dihydro-5H-benzo[b]pyrido[2,3-e][1,4]diazepine of Example 1, Step B (0.100 g, 0.51 mmol) in tetrahydrofuran (5 mL) was added 4-dimethylamino pyridine (0.190 g, 1.55 mmol) followed by 2-chloro-4-fluoro benzoylchloride (0.100 mL, 0.76 mmol). The mixture was stirred overnight at room temperature and then evaporated to dryness. The residue was partitioned between aqueous saturated ammonium chloride and dichloromethane. The organic layer was dried over sodium sulfate and evaporated to dryness to provide the title compound identical to the material described in Example 1, Step D.

25

30

EXAMPLE 24**[2-Chloro-4-(5-methyl-1H-[1,2,4]triazol-3-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-methanone**

5

Step A. 2-Chloro-terephthalamic acid methyl ester

A mixture of 2-chloro-4-cyano benzoic acid methyl ester (12.4 g, 63.4 mmol) and potassium carbonate (1.3 g, 9.4 mmol) in dimethylsulfoxide (40 mL) was treated dropwise under cooling with 30% hydrogen peroxide (7.6 mL). The mixture was allowed to warm to room temperature and stirred overnight. The solution was quenched with water and the resulting precipitate collected by filtration. The crude material was dissolved in dichloromethane and absorbed on a silica gel Merck-60 flash column. Elution with a dichloromethane-methanol gradient (from 98:2 to 90:10) provided the title compound (10 g, 74%) as a white solid, m.p. 154-156°C.

NMR (DMSO-d₆, 400 MHz): δ 3.87 (s, 3H), 7.67 (s, 1H, NH), 7.86-7.91 (m, 2H), 8.00-8.01 (m, 1H), 8.20 (s, 1H, NH)

MS (EI, m/z): 213 [M]⁺

Anal. Calc'd for C₉H₈ClNO₃: C 50.60, H 3.77, N 6.56. Found: C 50.36, H 3.66, N 6.44

20

Step B. 2-Chloro N-(1-dimethylaminoethylidene)-terephthalamic acid methyl ester

A mixture of 2-chloro-terephthalamic acid methyl ester of Step A (1.02 g, 4.8 mmol) and N,N -dimethylacetamide dimethyl acetal (3.5 mL, 23.9 mmol) was heated at 90°C for 30 minutes under nitrogen. The solution was cooled, and excess reagent was removed under high vacuum to provide a brown oil which was used as such in the next step.

NMR (DMSO-d₆, 400 MHz): δ 2.29 (s, 3H), 3.14 (s, 3H), 3.16 (s, 3H), 3.87 (s, 3H), 7.83-7.85 (m, 1H), 8.00-8.06 (m, 2H)

MS (EI, m/z): 282 [M]⁺

30

35

Step C. 2-Chloro-4-(5-methyl-1H-[1,2,4]triazol-3-yl)-benzoic acid methyl ester

Anhydrous hydrazine (0.30 mL, 9.6 mmol) was added via syringe to a solution of the intermediate of Step B (4.8 mmol) in glacial acetic acid (6 mL) under a nitrogen atmosphere. The reaction was heated at 90°C for 30 minutes, then cooled and concentrated in vacuo to a light brown solid. The solid was redissolved in aqueous methanol and the solution neutralized with saturated aqueous sodium bicarbonate. The mixture was extracted with dichloromethane and ethyl acetate, the extracts were combined and dried over sodium sulfate. Evaporation of the solvents yielded a solid which was triturated with ether to provide the title product (0.81 g, 67%) as an off-white solid, m.p. 196-198°C.

NMR (DMSO- d_6 , 400 MHz): δ 2.41 (s, 3H), 3.86 (s, 3H), 7.90-8.05 (m, 3H), 13.94 (s, 1H)

MS (EI, m/z): 251 [M]⁺

Anal. Calc'd for C₁₁H₁₀ClN₃O₂: C 52.50, H 4.01, N, 16.70. Found: C 52.68, H 3.83, N 16.50

Step D. 2-Chloro-4-[1-(4-methoxy-benzyl)-5-methyl-1H-[1,2,4]triazol-3-yl]-benzoic acid methyl ester 0.03 solvate with dichloromethane

Sodium hydride (60% suspension in oil, 0.30 g, 7.5 mmol) was washed with hexane and resuspended in dry dimethylformamide (20 mL) under a nitrogen atmosphere. The triazole intermediate of Step C (1.36 g, 5 mmol) was added and the mixture was stirred for one hour. p-Methoxybenzyl chloride (0.75 mL, 5.5 mmol) was added and after stirring for 3 hours, the reaction was quenched with water and extracted with ethyl acetate. The extracts were combined, dried over anhydrous sodium sulfate and concentrated in vacuo. The residue was dissolved in dichloromethane and absorbed on a silica Merck 60 flash column. Elution with 3% ethyl acetate in dichloromethane provided the title compound (1.23 g, 66.2%) as a white solid, m.p. 102-104°C.

-87-

NMR (DMSO- d_6 , 400 MHz): δ 2.48 (s, 3H), 3.72 (s, 3H), 3.86 (s, 3H), 5.35 (s, 2H), 6.90-6.92 (m, 2H), 7.23-7.25 (m, 2H), 7.89-8.02 (m, 3H)

MS (EI, m/z): 371 [M]⁺

Anal. Calc'd for $C_{19}H_{18}ClN_3O_3 + 0.03 CH_2Cl_2$: C 61.05, H 4.86, N 11.22. Found:

5 C 60.83, H 4.96, N, 11.18

Step E. 2-Chloro-4-[1-(4-methoxy-benzyl)-5-methyl-1H-[1,2,4]-triazol-3-yl]-benzoic acid 0.10 hydrate 0.04 solvate with ethyl acetate

10 A solution of the ester intermediate of Step D (1.6 g, 4.3 mmol) in methanol (15 mL) was treated with 2.5 N aqueous sodium hydroxide (3.5 mL, 8.8 mL) under a nitrogen atmosphere. The mixture was refluxed for two hours, cooled and concentrated in vacuo. The residue was partitioned between ethyl acetate and water. The aqueous layer was acidified with 1N aqueous HCl. The precipitate was collected
15 by filtration to provide the title compound (1.25 g, 81.2%) as a white solid, m.p. 154-156°C.

NMR (DMSO- d_6 , 400 MHz): δ 2.47 (s, 3H), 3.72 (s, 3H), 5.34 (s, 2H), 6.90-6.93 (m, 2H), 7.23-7.25 (m, 2H), 7.87-7.99 (m, 3H), 13.40 (s, 1H)

MS (EI, m/z): 357 [M]⁺

20 Anal. Calc'd for $C_{18}H_{16}ClN_3O_3 + 0.10 H_2O + 0.04 C_4H_8O_2$: C 60.07, H 4.59, N 11.57.

Found: C 59.75, H 4.41, N 11.43

Step F. 2-Chloro-4-[1-(4-methoxy-benzyl)-5-methyl-1H-[1,2,4]-triazol-3-yl]-benzoyl chloride
25

A suspension of the acid of Step E (1 g, 2.8 mmol) in dichloromethane containing a few drops of dimethylformamide was treated dropwise under nitrogen with oxalyl chloride (0.30 mL, 3.4 mmol). After gas evolution subsided, the reaction
30 mixture was refluxed for another 15 minutes and then evaporated to dryness in vacuo to provide the title compound which was used as such in the next step.

Step G. {2-Chloro-4-[1-(4-methoxy-benzyl)-5-methyl-1H-[1,2,4]triazol-3-yl]-phenyl}-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-methanone 0.06 dichloromethane solvate

- 5 To a solution of 6,11-dihydro-5H-pyrido[2,3-b][1,5]benzodiazepine of Example 1, Step B (0.55 g, 2.8 mmol) in dimethylformamide (10 mL) under nitrogen was added solid potassium carbonate (0.39 g, 2.8 mmol). The mixture was treated dropwise with a solution of the crude acid chloride (2.8 mmol) of Step F in dimethylformamide (10 mL). After stirring at room temperature for 90 minutes, the
- 10 reaction mixture was diluted with water and extracted with ethyl acetate. The organic extracts were combined and washed with 1N aqueous sodium hydroxide, dried over sodium sulfate and evaporated to dryness. The residue was dissolved in dichloromethane and absorbed on a column of flash silica Merck-60. Less polar impurities were eluted with 1:1 ethyl acetate-hexane. Further elution with 2%
- 15 methanol in dichloromethane provided the title compound as a white solid (0.57 g, 38%), m.p. 218-221°C.
- NMR (DMSO-d₆, 400 MHz): δ 2.42 (s, 3H), 3.71 (s, 3H), 4.14 and 5.44 (dd, 2H), 5.29 (s, 2H), 6.49 (m, 1H), 6.74-6.80 (m, 2H), 6.88-6.99 (m, 3H), 7.18-7.26 (m, 4H), 7.60 (m, 1H), 7.65-7.75 (m, 2H, ArH), 8.11 (m, 1H), 9.55 (s, 1H)
- 20 MS (ESI, m/z): 537 [M+H]⁺
- Anal. Calc'd for C₃₀H₂₅ClO₂ + 0.06 CH₂Cl₂ : C 66.60, H 4.67, N 15.50. Found: C 66.24, H 4.85, N 15.23

Step H. [2-Chloro-4-(5-methyl-1H-[1,2,4]-triazol-3-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-methanone

- 25 A solution of the triazole intermediate of Step G (0.54 g, 1.01 mmol) in trifluoroacetic acid (15 mL) was heated at reflux for seven days under a nitrogen atmosphere. The mixture was cooled and the trifluoroacetic acid removed in vacuo.
- 30 The residue was dissolved in water and neutralized with saturated aqueous sodium bicarbonate. The mixture was extracted with ethyl acetate, the extracts were dried over sodium sulfate and concentrated in vacuo to give a pale yellow solid. The residue was dissolved in ethyl acetate-methanol and absorbed on a silica Merck-60 flash column.

Elution with a solvent gradient (from 100% ethyl acetate to 5% methanol in ethyl acetate) provided the title compound (0.23 g, 54.6%) as a white solid, m. p. >270°C.

NMR (DMSO-d₆, 400 MHz): δ 2.36 (s, 3H), 4.15 and 5.45 (dd, 2H, CONCH₂), 6.50 (m, 1H), 6.75-6.80 (m, 2H), 6.98 (m, 1H), 7.19-7.27 (m, 2H), 7.60 (m, 1H), 7.70-7.79 (m, 2H), 8.11 (m, 1H), 9.54 (s, 1H), 13.78 (s, 1H)

MS (+FAB, m/z): 417 [M+H]⁺

Anal. Calc'd for C₂₂H₁₇ClN₆O: C 63.39, H 4.11, N 20.16. Found: C 63.14, H 4.13, N, 19.90

10

EXAMPLE 25

[2-Bromo-4-(3-methyl-pyrazol-1-yl)-phenyl]-(11H-5-oxa-4,10-diaza-dibenzo[a,d]cyclohepten-10-yl)-methanone

15

The title compound may be prepared in a manner analogous to that of the 2-chloro analog of Example 16, by reacting 6H-pyrido[2,3-b][1,5]benzoxazepine of Example 16, Step A, with 2-bromo-4-fluorobenzoyl chloride of Example 3, Step A. Subsequent reaction of the intermediate (2-bromo-4-fluoro-phenyl)-(11H-5-oxa-4,10-diaza-dibenzo[a,d] cyclohepten-10-yl)-methanone with the sodium salt of 3-methylpyrazole in a manner analogous to that of Example 16, Step C will provide the title compound.

20

EXAMPLE 26

25

[4-(3-Methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-(11H-5-oxa-4,10-diaza-dibenzo[a,d]cyclohepten-10-yl)-methanone

The title compound may be prepared in a manner analogous to that of the 2-chloro analog of Example 16, by reacting 6H-pyrido[2,3-b][1,5]benzoxazepine of Example 16, Step A with 2-trifluoromethyl-4-fluorobenzoyl chloride of Example 4, Step A. Subsequent reaction of the intermediate (4-fluoro-2-trifluoromethyl-phenyl)-(11H-5-oxa-4,10-diaza-dibenzo[a,d]cyclohepten-10-yl)-methanone with the sodium salt of 3-methylpyrazole in a manner analogous to that of Example 16, Step C will provide the title compound.

35

EXAMPLE 27**[2-Fluoro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(11H-5-oxa-4,10-diaza-dibenzo[a,d]cyclohepten-10-yl)-methanone**

5

The title compound may be prepared in a manner analogous to that of the 2-chloro analog of Example 16, by reacting 6H-pyrido[2,3-b][1,5]benzoxazepine of Example 16, Step A with 2,4-difluoro benzoylchloride of Example 9, Step A. Subsequent reaction of the intermediate (2,4-difluoro-phenyl)-(11H-5-oxa-4,10-diaza-dibenzo[a,d]cyclohepten-10-yl)-methanone with the sodium salt of 3-methylpyrazole in a manner analogous to that of Example 16, Step C will provide the title compound.

15


EXAMPLE 28**[2-Chloro-4-(1-methyl-1H-pyrazol-3-yl)-phenyl]-(11H-5-oxa-4,10-diaza-dibenzo[a,d]cyclohepten-10-yl)-methanone**

20

The title compound may be prepared in a manner analogous to that of the pyrazole analog of Example 19, by reacting the intermediate 2-chloro-4-(1-methyl-1H-pyrazol-3-yl)-benzoylchloride of Example 19 with 6H-pyrido[2,3-b][1,5]benzoxazepine of Example 16, Step A.

CLAIMS

1. A compound of the general formula (I):




(I)

W is O, or NR⁶

A and B are, independently, carbon or nitrogen;

R¹ is $\text{—C}\equiv\text{C—R}^9$, alkanoyl of 2 to 7 carbon atoms or a moiety selected from the group of:



R^2 , R^3 and R^5 are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or perfluoroalkyl of 1 to 6 carbons;

R^4 is hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, alkoxyalkyl of 2 to 7 carbon atoms, optionally substituted aralkyl of 7 to 12 carbon atoms, or an acyl substituent selected from the group consisting of alkanoyl of 2 to 7 carbon atoms, alkenoyl of 3 to 7 carbon atoms, cycloalkanoyl of 3 to 7 carbon atoms, arylalkanoyl having an alkane chain of from 1 to 6 carbon atoms, aroyl or heteroaroyl of 7 to 13 carbon atoms;

R^6 is hydrogen, acyl of 2 to 6 carbon atoms, straight chain alkyl of 1 to 6 carbon atoms, or branched chain alkyl of 3 to 7 carbon atoms;

X and Y are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, alkoxyalkyl of 2 to 7 carbon atoms, halogen, straight or branched chain alkoxy of 1 to 6 carbon atoms, hydroxy, CF_3 , or perfluoroalkyl of 2 to 6 carbons;

Z is hydrogen or a straight chain alkyl group of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, halogen, alkoxyalkyl of 2 to 7 carbons, or hydroxyalkyl of 1 to 6 carbons, or $\text{CH}_2\text{NR}^7\text{R}^8$;


R^7 and R^8 are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, aryl, or arylalkyl; or taken together with the nitrogen they form a five or six membered ring optionally containing one or more additional heteroatoms;

R^9 is independently, hydrogen, a silyl containing group, or a lower alkyl of 1 to 6 carbons; and

R^{10} is a straight chain alkyl of 1 to 6 carbon atoms;

or a pharmaceutically acceptable salt thereof.


2. A compound as claimed in Claim 1 wherein R^7 and R^8 taken together are a five or six membered ring selected from the group:



or a pharmaceutically acceptable salt thereof.


3. A compound selected from Formula (Ia):

-94-



wherein:

R^1 is alkanoyl of 2 to 7 carbon atoms or a group selected from



R^2 , R^3 and R^5 are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or perfluoroalkyl of 1 to 6 carbons;

R^4 is hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, alkoxyalkyl of 2 to 7 carbon atoms, optionally substituted aralkyl of 7 to 12 carbon atoms, or an acyl substituent selected from the group consisting of alkanoyl of 2 to 7 carbon atoms, alkenoyl of 3 to 7 carbon atoms, cycloalkanoyl of 3 to 7 carbon atoms, aroyl or

heteroaroyl of 7 to 12 carbon atoms, optionally substituted with one or two alkyl groups of 1 to 6 carbon atoms, or arylalkanoyl of 1 to 6 carbon atoms;

R^6 is hydrogen, acyl of 2 to 6 carbon atoms, straight chain alkyl of 1 to 6 carbon atoms, or branched chain alkyl of 3 to 7 carbon atoms;

X and Y are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, alkoxyalkyl of 2 to 7 carbon atoms, halogen (including chlorine, bromine, fluorine, and iodine), straight or branched chain alkoxy of 1 to 6 carbons, hydroxy, CF_3 , or perfluoroalkyl of 2 to 6 carbons;


Z is hydrogen or a straight chain alkyl group of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, halogen, alkoxyalkyl of 2 to 7 carbons, or hydroxyalkyl of 1 to 6 carbons, or $CH_2NR^7R^8$;

R^7 and R^8 are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms or taken together with the nitrogen they form a five or six membered ring optionally containing one or more additional heteroatoms; and

R^{10} is a straight chain alkyl of 1 to 6 carbon atoms;


or a pharmaceutically acceptable salt thereof.

4. A compound as claimed in Claim 3 wherein R^7 and R^8 taken together are a five or six membered ring selected from the group:




or a pharmaceutically acceptable salt thereof.

5. A compound selected from Formula (Ib):



wherein:

R^1 is alkanoyl of 2 to 7 carbon atoms or a group selected from



R^2 , R^3 and R^5 are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, or perfluoroalkyl of 1 to 6 carbons;

R^4 is hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, alkoxyalkyl of 2 to 7

carbon atoms, optionally substituted aralkyl of 7 to 12 carbon atoms, or an acyl substituent selected from the group consisting of alkanoyl of 2 to 7 carbon atoms, alkenoyl of 3 to 7 carbon atoms, cycloalkanoyl of 3 to 7 carbon atoms, aroyl or heteroaroyl of 7 to 12 carbon atoms, optionally substituted with one or two alkyl groups of 1 to 6 carbon atoms, or arylalkanoyl of 1 to 6 carbon atoms;

R^6 is hydrogen, acyl of 2 to 6 carbon atoms, straight chain alkyl of 1 to 6 carbon atoms, or branched chain alkyl of 3 to 7 carbon atoms;

X and Y are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, alkoxyalkyl of 2 to 7 carbon atoms, halogen (including chlorine, bromine, fluorine, and iodine), straight or branched chain alkoxy of 1 to 6 carbons, hydroxy, CF_3 , or perfluoroalkyl of 2 to 6 carbons;

Z is hydrogen or a straight chain alkyl group of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms, cycloalkyl of 3 to 7 carbon atoms, halogen, alkoxyalkyl of 2 to 7 carbons, or hydroxyalkyl of 1 to 6 carbons, or $CH_2NR^7R^8$;


R^7 and R^8 are, independently, hydrogen, straight chain alkyl of 1 to 6 carbon atoms, branched chain alkyl of 3 to 7 carbon atoms or taken together with the nitrogen they form a five or six membered ring optionally containing one or more additional heteroatoms; and

R^{10} is a straight chain alkyl of 1 to 6 carbon atoms;

or a pharmaceutically acceptable salt thereof.

6. A compound as claimed in Claim 5 wherein R^7 and R^8 taken together are a five or six membered ring selected from the group:

-98-



or a pharmaceutically acceptable salt thereof.

7. A compound as claimed in Claim 1 which is [2-Chloro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-methanone,

[[2-Chloro-4-(5-methyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-methanone,

[[2-Bromo-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone,

[(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-(4-fluoro-2-trifluoromethyl-phenyl)-methanone,

[(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[4-(3-methyl-pyrazol-1-yl) -2-trifluoromethyl-phenyl]-methanone,

[(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[4-(3-methyl-pyrazol-1-yl) -2-trifluoromethyl-phenyl]-methanone 1:1 salt with methanesulfonic acid,

[(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[4-(3-methyl-pyrazol-1-yl) -2-trifluoromethyl-phenyl]-methanone 1:1 salt with hydrochloric acid,

[4-(3-Methyl-pyrazol-1-yl)-2-trifluoromethyl-benzoic acid methylester,

[4-(3-Methyl-pyrazol-1-yl)-2-trifluoromethyl-benzoic acid or a pharmaceutically acceptable salt or ester thereof,

[(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[4-(5-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-methanone,

[(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-[2-trifluoromethyl-4-(3-trifluoromethyl-pyrazol-1-yl)-phenyl]-methanone,

[(5-Methyl-5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[4-(3-methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-methanone,

[(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[2-fluoro-4-(3-methyl-pyrazol-1-yl)-phenyl]-methanone,

[(2,4-Difluoro-phenyl)-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone,

[(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[4-fluoro-2-(3-methyl-pyrazol-1-yl)-phenyl]-methanone,

[[2-Chloro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5-methyl-5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone,

[(2-Chloro-4-fluoro-phenyl)-(5-methyl-5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone,

[(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-[2-methyl-5-(3-methyl-pyrazol-1-yl)-phenyl]-methanone,

[(5,11-Dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-(5-fluoro-2-methyl-phenyl) -methanone,

[[4-(3-tert-Butyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone,

[[2-Chloro-4-(3-trifluoromethyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-methanone,

[[2-Chloro-4-(1-methyl-1H-pyrazol-3-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone,

[[2-Chloro-4-(1-methyl-1H-pyrazol-3-yl)-phenyl]-(5-methyl-5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-10-yl)-methanone,

[6,11-Dihydro-5H-pyrido-[2,3-b][1,5]-benzodiazepine-5-one,

[[2-Chloro-4-(5-methyl-1H-[1,2,4]triazol-3-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-methanone,

[2-Chloro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(5,11-dihydro-pyrido[2,3-b][1,5]benzodiazepin-6-yl)-methanone,

[2-Bromo-4-(3-methyl-pyrazol-1-yl)-phenyl]-(11H-5-oxa-4,10-diaza-dibenzo[a,d]cyclohepten-10-yl)-methanone,

[4-(3-Methyl-pyrazol-1-yl)-2-trifluoromethyl-phenyl]-(11H-5-oxa-4,10-diaza-dibenzo[a,d]cyclohepten-10-yl)-methanone,

[2-Fluoro-4-(3-methyl-pyrazol-1-yl)-phenyl]-(11H-5-oxa-4,10-diaza-dibenzo[a,d]cyclohepten-10-yl)-methanone,

or [2-Chloro-4-(1-methyl-1H-pyrazol-3-yl)-phenyl]-(11H-5-oxa-4,10-diaza-dibenzo[a,d]cyclohepten-10-yl)-methanone.

8. A method of treating disorders in a mammal which are remedied or alleviated by vasopressin agonist activity, the method comprising administering to the mammal in need thereof a pharmaceutically effective amount of a compound as claimed in any one of Claims 1 to 7, or a pharmaceutically acceptable salt, ester or prodrug form thereof, and a suitable pharmaceutical carrier.

9. A method as claimed in Claim 8 wherein the disorder which is remedied or alleviated by vasopressin agonist activity is selected from the group of diabetes

insipidus, nocturnal enuresis, nocturia, urinary incontinence, bleeding and coagulation disorders, or the inability to temporarily delay urination.

10. A pharmaceutical composition useful for treating disorders which are remedied or alleviated by vasopressin agonist activity in a mammal, the composition comprising a pharmaceutically effective amount of a compound as claimed in any one of Claims 1 to 7 in combination or association with a pharmaceutically acceptable carrier.


11. A pharmaceutical composition as claimed in Claim 10 wherein the disorder remedied or alleviated by vasopressin agonist activity in a mammal is selected from the group of diabetes insipidus, nocturnal enuresis, nocturia, urinary incontinence, bleeding and coagulation disorders, or the inability to temporarily delay urination.

12. Use of a compound as claimed in any one of claims 1 to 7 as a medicament.

13. Use of a compound as claimed in any one of claims 1 to 7 in the preparation of a medicament for the treatment of diabetes insipidus, nocturnal enuresis, nocturia, urinary incontinence, bleeding and coagulation disorders, or the inability to temporarily delay urination.

14. A process for the preparation of a compound of formula I which comprises:

a) reacting a compound of formula




wherein W, A, B, X, Y and Z are each as defined Claim 1,

with the appropriate compound of formula R^1H , wherein R^1 is a heterocyclic moiety selected from groups a, b, c, d, l, n and o defined Claim 1,


to provide the desired compound of formula I wherein R^1 is the a heterocyclic moiety selected from groups a, b, c, d, l, n and o defined Claim 1,

b) reacting a compound of formula



wherein W, X and Z are each as defined Claim 1,


with an acylating agent of formula 9



wherein A and B are as defined Claim 1, J is an acylating moiety and R^1 is a heterocyclic moiety selected from groups a, b, c, d, l, n and o defined Claim 1;

to provide the desired compound of formula I wherein R^1 is a heterocyclic moiety selected from groups a, b, c, d, l, n and o defined Claim 1;


c) reacting a compound of formula



wherein W is oxygen or NH, and X and Z are as defined Claim 1,

-103-


with a compound of formula



wherein A and B are both carbon, J is an acylating moiety and R¹ is a heterocyclic moiety selected from groups g, f and h defined Claim 1,

to provide a compound of formula I wherein W is oxygen or NH, A and B are both carbon and R¹ is a heterocyclic moiety selected from groups g, f and h defined Claim 1;

d) reacting a compound of formula 40




wherein W, A, B, X, Z and R² are each as defined Claim 1,

with hydroxylamine or an appropriate substituted hydrazine of formula R⁴-NHNH₂ wherein R⁴ is as defined Claim 1,

to provide the desired compound of formula I wherein W, A, B, X, Y and Z are each as defined Claim 1 and R¹ is a heterocyclic moiety selected from groups f, g and j defined Claim 1;

e) reacting a compound of formula 48

-104-




wherein W is oxygen or NH, A, B, X, Y and Z are each as defined Claim 1,

with hydroxylamine or an appropriate substituted hydrazine of formula $R^4\text{-NHNH}_2$ wherein R^4 is as defined Claim 1


to provide the desired compound of formula I wherein W is oxygen or NH, A, B, X, Y and Z are each as defined Claim 1 and R^1 is a heterocyclic moiety selected from groups e, i and k defined Claim 1;

f) reacting a compound of formula



wherein W is oxygen or NH and X and Z are each as defined Claim 1,

with a compound of formula




wherein J is an acylating moiety, R^1 is a heterocyclic moiety selected from groups e and i defined Claim 1 and R^4 is other than hydrogen,

to provide a compound of formula I wherein W is oxygen or NH, R¹ is a heterocyclic moiety selected from groups e and i defined Claim 1 and R⁴ is other than hydrogen;

g) converting a compound of formula I wherein W is oxygen or NH, A, B, X, Y, Z and R² are each as defined Claim 1, R¹ is an heterocyclic moiety selected from groups e and i defined Claim 1 and R⁴ is an optionally substituted aralkyl group

to the corresponding compound of formula I wherein W is oxygen or NH, A, B, X, Y, Z and R² are each as defined Claim 1, R¹ is an heterocyclic moiety selected from groups e and i defined Claim 1 and R⁴ is hydrogen;

h) converting a compound of formula 54



(54)

wherein W, A, B, X, Y and Z are each as defined Claim 1


to the desired compound of formula I wherein W, A, B, X, Y and Z are each as defined Claim 1 and R¹ is heterocyclic moiety m defined Claim 1;

i) alkylating or acylating a compound of formula I wherein W is NR⁶, R⁶ is hydrogen, X, Y and Z are each as defined Claim 1, A and B are both C and R¹ is heterocyclic moiety selected from groups e, f, g, h, i, j, and k defined Claim 1,

to provide the corresponding compound of formula I wherein W is NR⁶ and R⁶ is other than hydrogen.

j) converting a compound of formula 63


-106-



wherein A, B, X and Z are each as defined Claim 1 and R⁶ is other than hydrogen,

to the corresponding compound of formula I wherein A, B, X and Z are each as defined Claim 1, W is NR⁶, R⁶ is other than hydrogen and R¹ is heterocyclic moiety m defined Claim 1;

k) converting a compound of formula 64




wherein A, B, X and Z are each as defined Claim 1 and R⁶ is other than hydrogen

to the corresponding compound of formula I wherein A, B, X and Z are each as defined Claim 1, W is NR⁶, R⁶ is other than hydrogen and R¹ is a heterocyclic moiety selected from groups e, i and k defined Claim 1;

l) reacting a compound of formula 1


-107-



(1)

wherein W, X and Z are as defined in Claim 1,


with a compound of formula



wherein A, B and Y are as defined in Claim 1, J is an acylating group and R¹⁰ is an alkyl group,

to provide a compound of formula I wherein W, A, B, X and Z are each as defined Claim 1 and R¹ is a an alkanoyl group;


m) reacting a compound of formula



wherein W, A, B, X and Z are each as defined Claim 1

with a compound of formula 53

-108-



wherein R^9 is as defined Claim 1

to provide a compound of formula I wherein W, A, B, X and Z are each as defined Claim 1, R^1 is $\text{—C}\equiv\text{C}-\text{R}^9$ and R^9 is as defined Claim 1

or n) converting a compound of formula I which is a free base to the corresponding pharmaceutically acceptable salt.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/15487

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D471/04 C07D498/04 A61K31/55

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 512 563 A (DU XUEMEI ET AL) 30 April 1996	1-6,8-14
X	see claim 1	7
A	WO 96 22282 A (AMERICAN CYANAMID CO) 25 July 1996	1-6,8-14
X	see claim 1	7
A	US 5 420 270 A (CHANDRAKUMAR NIZAL S ET AL) 30 May 1995	1-6,8-14
X	see formula I	7
A	FR 4 999 M (KARL THOMAE GMBH) 17 April 1967	1-6,8-14
X	see formula I and page 2, right hand column	7
-/--		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

2 October 1998

Date of mailing of the international search report

12/10/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Steendijk, M

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 98/15487

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 97 22591 A (OTSUKA PHARMA CO LTD ;OGAWA HIDENORI (JP); KONDO KAZUMI (JP); SHIN) 26 June 1997 see claim 1</p> <p>-----</p>	1-14

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/US 98/15487

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5512563 A	30-04-1996	US 5739128 A	14-04-1998
		US 5686445 A	11-11-1997
		US 5760031 A	02-06-1998
		US 5736538 A	07-04-1998
		US 5747487 A	05-05-1998
		US 5693635 A	02-12-1997
		US 5786353 A	28-07-1998
		AU 676737 B	20-03-1997
		AU 6877694 A	09-02-1995
		CA 2128955 A	30-01-1995
		CN 1106802 A	16-08-1995
		CZ 9401798 A	15-02-1995
		EP 0640592 A	01-03-1995
		FI 943542 A	30-01-1995
		HU 71548 A	28-12-1995
		JP 7179430 A	18-07-1995
		NO 942817 A	30-01-1995
		NZ 264116 A	26-11-1996
		PL 304496 A	06-02-1995
		SK 88094 A	10-05-1995
		ZA 9405604 A	09-03-1995
WO 9622282 A	25-07-1996	AU 4904296 A	07-08-1996
		BR 9606977 A	04-11-1997
		CA 2210688 A	25-07-1996
		CZ 9702245 A	17-12-1997
		EP 0804420 A	05-11-1997
US 5420270 A	30-05-1995	US 5512561 A	30-04-1996
FR 4999 M	17-04-1967	CH 453374 A	
		DE 1251767 B	
		FR 1458125 A	25-01-1967
		GB 1071757 A	
		US 3326900 A	20-06-1967
WO 9722591 A	26-06-1997	AU 1110997 A	14-07-1997
		JP 10081668 A	31-03-1998